

CHARACTER SHEAVES ON DISCONNECTED GROUPS, VIII

G. LUSZTIG

INTRODUCTION

Throughout this paper, G denotes a fixed, not necessarily connected, reductive algebraic group over an algebraically closed field $\mathbf{k}$. This paper is a part of a series [L9] which attempts to develop a theory of character sheaves on G .

In Section 36 we associate to any subset J of the set of simple reflections an algebra $\underline{\mathbf{K}}^J$ over $\mathbf{Q}(v)$ (with v an indeterminate) defined using certain character sheaves on Z_{J,G^0} . When $J = \emptyset$ this is the standard Iwahori-Hecke algebra attached to $\mathbf{W}$. For general J we show that this algebra shares several basic features with an Iwahori-Hecke algebra with unequal parameters. This opens the possibility of studying Iwahori-Hecke algebras with unequal parameters in the framework of the theory of perverse sheaves.

In Section 37 we prove a Mackey type formula for character sheaves on $Z_{J,D}$ where $J \subset \mathbf{I}$ and D is a connected component of G . This is essentially an identity involving certain induction and restriction functors analogous to one in the representation theory of reductive groups over a finite field.

In Section 38 we study a duality operation for character sheaves on a connected component D of G generalizing the case $D = G^0$ considered in [L3, III, 15]. This duality operator is analogous to the known duality operator for representations of a reductive group over a finite field, see 38.12.

In Section 39 we prove a quasi-rationality property for representations of certain extensions of an (irreducible) Weyl group. This generalizes [L3, III, (12.9.3)] and is a step in the proof of a key property of character sheaves (see 39.8).

Errata to Part V.

In 27.1 replace the diagram $N_DP/U_P \xleftarrow{a} V_1 \xrightarrow{a} V_2 \xrightarrow{a'} D$ by $N_DP/U_P \xleftarrow{a} V_1 \xrightarrow{a'} V_2 \xrightarrow{a''} D$.

In the commutative diagram on p.372 replace $\hat{Z}_1 \xrightarrow{a} \hat{Z}_2, Z_1 \xrightarrow{b} Z_2, V_1 \xrightarrow{c} V_2 \xrightarrow{a'} D$ by $\hat{Z}_1 \xrightarrow{a'} \hat{Z}_2, Z_1 \xrightarrow{b'} Z_2, V_1 \xrightarrow{c'} V_2 \xrightarrow{a''} D$ respectively.

Errata to Part VI.

In 28.5 replace $t_0 : \text{Ad}(d^{-1}\dot{w}^{-1})(t_0)tt_0^{-1}$ by $t_0 : t \mapsto \text{Ad}(d^{-1}\dot{w}^{-1})(t_0)tt_0^{-1}$.

Supported in part by the National Science Foundation.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

In 28.19(a) replace $\hat{Z}_{\emptyset, \epsilon_D(J), D^{-1}}^{\mathcal{L}'}$ by $\hat{Z}_{\epsilon_D(J), D^{-1}}^{\mathcal{L}'}$ (twice).

Errata to Part VII.

In 32.5 replace $> b_2$ by b_2 .

In 32.15 replace C_u by C^u .

In 32.18(a) replace $T_{\epsilon'(a_{r+r'}^{-1})}$ by $T_{\epsilon'(a_{r+r'}^{-1})}$.

In 32.23 replace the first $=$ by $) =$.

After the statement of Corollary 32.23 insert: Take $D' = D^{-1}$ so that $\Delta = G^0$.

In the third line of Corollary 32.24 replace λ' by λ'' .

In the fourth line of Corollary 32.24 replace λ'' by $\mathcal{L}''$.

In 32.26, line 13, replace tT_x by $\tilde{T}_x$.

In 35.6, line 8, after $h_{i-1}^{-1}h_i \in U^*n_iU^*$, add: ($i \in [1, r]$).

CONTENTS

36. The algebra $\underline{\mathfrak{K}}^J$.

37. A Mackey type formula.

38. Duality.

39. Quasi-rationality.

36. THE ALGEBRA $\underline{\mathfrak{K}}^J$

36.1. We fix a connected component D of G . We write ϵ instead of $\epsilon_D : \mathbf{W} \rightarrow \mathbf{W}$ (see 26.2). Occassionally we shall consider one (or two) other component(s), say D' (or D', D''); we write ϵ' instead of $\epsilon_{D'}$.

For $J \subset \mathbf{I}$ we identify $Z_{J,D}$ (see 26.2) with

$$\{(P, gU_P); P \in \mathcal{P}_J, gU_P \in D/U_P\}$$

by $(P, P', gU_P) \mapsto (P, gU_P)$. To any $(P, gU_P) \in Z_{J,D}$ we associate an element $w_{P, gU_P} \in \mathbf{W}$ by the following requirements. Let $z = \text{pos}(gPg^{-1}, P) \in {}^{\epsilon(J)}\mathbf{W}^J$; then

(i) $w_{P, gU_P} = w_{P_1, gU_{P_1}}$ where $P_1 = (g^{-1}Pg \cap P)U_P \in \mathcal{P}_{J \cap \epsilon^{-1}(\text{Ad}(z)J)}$,

(ii) $w_{P, gU_P} = z$ if $\epsilon^{-1}(\text{Ad}(z)J) = J$.

Note that (i),(ii) define uniquely w_{P, gU_P} by induction on $|J|$: if $|J| = 0$ then w_{P, gU_P} is given by (ii); if $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(z)J) = J$ then w_{P, gU_P} is again given by (ii); if $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(z)J) \neq J$ then $|J \cap \epsilon^{-1}(\text{Ad}(z)J)| < |J|$ and w_{P, gU_P} is determined by (i) since $w_{P_1, gU_{P_1}}$ is known from the induction hypothesis.

From definitions we see that the map $Z_{J,D} \rightarrow \mathbf{W}$, $(P, gU_P) \mapsto w_{P, gU_P}$ is the composition of $\beta' : Z_{J,D} \rightarrow \mathcal{T}(J, \epsilon)$ in 26.2 (see also [L10, 3.11]) with the bijection $\mathcal{T}(J, \epsilon) \rightarrow {}^{\epsilon(J)}\mathbf{W}$ given by [L10, 2.4, 2.5] and with the inclusion ${}^{\epsilon(J)}\mathbf{W} \rightarrow \mathbf{W}$. In particular, we have $w_{P, gU_P} \in {}^{\epsilon(J)}\mathbf{W}$ and $\text{pos}(gPg^{-1}, P) = \min(\mathbf{W}_{\epsilon(J)} w_{P, gU_P} \mathbf{W}_J)$ for any $(P, gU_P) \in Z_{J,D}$. (For any $\mathbf{W}_{\epsilon(J)}, \mathbf{W}_J$ double coset Ω in $\mathbf{W}$ we denote by $\min(\Omega)$ the unique element of minimal length in Ω .)

For any $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$, we set

$${}^wZ_{J,D} = \{(P, gU_P) \in Z_{J,D}; w_{P,gU_P} = w\}.$$

Then $Z_{J,D} = \cup_{w \in {}^{\epsilon(J)}\mathbf{W}} {}^wZ_{J,D}$ is a partition; it is the same as the partition

$$Z_{J,D} = \cup_{\mathbf{t} \in \mathcal{T}(J,\epsilon)} {}^{\mathbf{t}}Z_{J,D}$$

in 26.2 (see also [L10, 3.11]). In particular, each ${}^wZ_{J,D}$ is a locally closed, smooth subvariety of $Z_{J,D}$ stable under the G^0 -action $h : (P, gU_P) \mapsto (hPh^{-1}, hgh^{-1}U_{hPh^{-1}})$. ■

From definitions, for $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$ we have a map

$$\vartheta_{J,w} : {}^wZ_{J,D} \rightarrow {}^wZ_{J_1,D}, (P, gU_P) \mapsto (P_1, gU_{P_1})$$

with $J_1 = J \cap \epsilon^{-1}(\text{Ad}(w_0)J)$, $w_0 = \min(\mathbf{W}_{\epsilon(J)} w \mathbf{W}_J)$, $P_1 = (g^{-1}Pg \cap P)U_P$. This map may be identified with a map ϑ as in [L10, 3.11]; in particular it is an affine space bundle (see [L10, 3.12(b)]).

36.2. If $J \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$ satisfies $\epsilon^{-1}(\text{Ad}(w)J) = J$ (hence $w \in {}^{\epsilon(J)}\mathbf{W}^J$), we have ${}^wZ_{J,D} = \{(P, gU_P) \in Z_{J,D}, \text{pos}(gPg^{-1}, P) = w\}$. In this case we pick $P \in \mathcal{P}_J, P' \in \mathcal{P}_{\epsilon(J)}$ such that $\text{pos}(P', P) = w$ and a common Levi L of P', P . Let $\mathbf{d} = \{g \in D; gLg^{-1} = L, gPg^{-1} = P'\}$, a connected component of the reductive group N_GL with identity component L . We have a diagram

$$(a) \quad \mathbf{d} \xleftarrow{pr_2} G^0 \times \mathbf{d} \xrightarrow{j} {}^wZ_{J,D}$$

where $j(h, g) = (hPh^{-1}, hgh^{-1}U_{hPh^{-1}})$ is a principal $(P \cap P')$ -bundle for a $(P \cap P')$ -action on $G^0 \times \mathbf{d}$ compatible under pr_2 with the $(P \cap P')$ -action on $\mathbf{d}$ via its quotient $(P \cap P')/U_{P \cap P'} = L$ (by conjugation in N_GL). If X is a character sheaf on $\mathbf{d}$ then $pr_2^\star X$ is therefore a $(P \cap P')$ -equivariant simple perverse sheaf on $G^0 \times \mathbf{d}$ which must be of the form $j^\star X'$ for a well defined simple perverse sheaf X' on ${}^wZ_{J,D}$.

The collection of simple perverse sheaves on ${}^wZ_{J,D}$ of the form X' with X as above is denoted by ${}^w\hat{Z}_{J,D}$. This collection is independent of the choice of P, P', L . Note that $X \mapsto X'$ defines a bijection between the set of isomorphism classes of character sheaves on $\mathbf{d}$ and the set of isomorphism classes of objects in ${}^w\hat{Z}_{J,D}$.

36.3. More generally, for $J \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, we define by induction on $|J|$ a collection of simple G^0 -equivariant perverse sheaves ${}^w\hat{Z}_{J,D}$ on ${}^wZ_{J,D}$.

If $|J| = 0$ then ${}^w\hat{Z}_{J,D}$ is defined as in 36.2. If $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(w)J) = J$, then ${}^w\hat{Z}_{J,D}$ is again defined as in 36.2. If $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(w)J) \neq J$ then $\epsilon^{-1}(\text{Ad}(w_0)J) \neq J$ where $w_0 = \min(\mathbf{W}_{\epsilon(J)} w \mathbf{W}_J)$. Thus, if $J_1 = J \cap \epsilon^{-1}(\text{Ad}(w_0)J)$ then $|J_1| < |J|$ and the class of perverse sheaves ${}^w\hat{Z}_{J_1,D}$ on ${}^wZ_{J_1,D}$ is defined from the induction hypothesis. By definition, ${}^w\hat{Z}_{J,D}$ consists of the simple perverse sheaves on ${}^wZ_{J,D}$ of the form $\vartheta_{J,w}^\star(X)$ for some $X \in {}^wZ_{J_1,D}$ (with $\vartheta_{J,w}$ as in 36.1). This completes the inductive definition of ${}^w\hat{Z}_{J,D}$. The objects of ${}^w\hat{Z}_{J,D}$ are

said to be *character sheaves* on ${}^wZ_{J,D}$. Let $\mathcal{I}_{J,w,D}$ be a set of representatives for the isomorphism classes of character sheaves on ${}^wZ_{J,D}$.

For $J \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, let $\mathcal{D}^{cs}({}^wZ_{J,D})$ be the subcategory of $\mathcal{D}({}^wZ_{J,D})$ whose objects are those $K \in \mathcal{D}({}^wZ_{J,D})$ such that for any j , any simple subquotient of ${}^pH^j K$ is in ${}^w\hat{Z}_{J,D}$. Let $i_{J,w} : {}^wZ_{J,D} \rightarrow Z_{J,D}$ be the inclusion.

(a) If $K \in \mathcal{D}^{cs}(Z_{J,D})$ then $i_{J,w}^*(K) \in \mathcal{D}^{cs}({}^wZ_{J,D})$.

It is enough to prove (a) for $K \in \hat{Z}_{J,D}$. In this case (a) follows from [L10, 4.12].

For $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$ and $K \in {}^w\hat{Z}_{J,D}$, let $K^\sharp$ be the unique simple perverse sheaf on $Z_{J,D}$ such that $i_{J,w}^*(K^\sharp) = K$ and $\text{supp}(K^\sharp)$ is the closure in $Z_{J,D}$ of $\text{supp}(K)$.

(b) We have $K^\sharp \in \hat{Z}_{J,D}$.

This is proved in [L10, 4.17] based on the following statement:

(c) Let Y be a locally closed subvariety of an algebraic variety Y' and let $i : Y \rightarrow Y'$ be the inclusion. Let $C \in \mathcal{D}(Y)$ and let A be a simple perverse sheaf on Y such that $A \dashv C$. Let $A^\sharp$ be the unique simple perverse sheaf on Y such that $i^*(A^\sharp) = A$ and $\text{supp}(A^\sharp)$ is the closure in Y' of $\text{supp}(A)$. Then $A^\sharp \dashv i_! C$.

(In *loc.cit.* this is applied with $Y = {}^wZ_{J,D}$, $Y' = Z_{J,D}$, $A = K$ and an $C \in \mathcal{D}^{cs}(Z_{J,D})$.) Since the proof of (c) is omitted in *loc.cit.* we give a proof here. Let

Y_1 be the closure of Y in Y' . Let $Y \xrightarrow{i_1} Y_1 \xrightarrow{i_2} Y'$ be the inclusions. Clearly, if (c) holds for i_1 and i_2 instead of i then it holds for i . Thus we may assume that $i = i_1$ or $i = i_2$ that is, that Y is open or closed in Y' . Assume first that Y is closed in Y' . Then $A^\sharp = i_! A$ and $i_!$ commutes with taking ${}^pH^j$; hence $i_! A \dashv i_! C$ as desired. Next, assume that Y is open in Y' . Then i^* commutes with taking ${}^pH^j$ and $i^* i_! C = C$. Hence for any j we have ${}^pH^j(C) = i^*({}^pH^j(i_! C))$ and for some j we have $A \dashv i^*({}^pH^j(i_! C))$. Let

$$0 = F_0 \subset F_1 \subset \dots \subset F_m = {}^pH^j(i_! C)$$

be a sequence of perverse subsheaves of ${}^pH^j(i_! C)$ such that F_k/F_{k-1} is simple for $k \in [1, m]$. Then

$$0 = i^* F_0 \subset i^* F_1 \subset \dots \subset i^* F_m = i^*({}^pH^j(i_! C))$$

is a sequence of perverse subsheaves of $i^*({}^pH^j(i_! C))$ such that $i^* F_k/i^* F_{k-1} = i^*(F_k/F_{k-1})$ is simple or 0 for $k \in [1, m]$. Hence $A \cong i^* F_k/i^* F_{k-1} = i^*(F_k/F_{k-1})$ for some $k \in [1, m]$. Then $A' = F_k/F_{k-1}$ is a simple perverse sheaf on Y' such that $A' \dashv i_! C$, $i^* A' \cong A$. We must have $A' \cong A^\sharp$ and (c) is proved.

(d) Let $K' \in \hat{Z}_{J,D}$. There exists a unique $w \in {}^{\epsilon(J)}\mathbf{W}$ and a unique $K \in {}^w\hat{Z}_{J,D}$ (up to isomorphism) such that $K' \cong K^\sharp$.

This is proved in [L10, 4.13]; it is an immediate consequence of (a).

Let $\mathcal{I}_{J,D} = \sqcup_{w \in {}^{\epsilon(J)}\mathbf{W}} \{K^\sharp; K \in \mathcal{I}_{J,w,D}\}$. From (b), (d) we see that

(e) $\mathcal{I}_{J,D}$ is a set of representatives for the isomorphism classes of character sheaves on $Z_{J,D}$.

From (a),(b) we deduce:

(f) If $w \in {}^{\epsilon(J)}\mathbf{W}$ and $K \in \mathcal{D}^{cs}({}^wZ_{J,D})$, then $(i_{J,w})_! K \in \mathcal{D}^{cs}(Z_{J,D})$.

36.4. For $J \subset J' \subset \mathbf{I}$ and $P \in \mathcal{P}_J$ let $Q_{J',P}$ be the unique parabolic in $\mathcal{P}_{J'}$ that

contains P . We have a diagram

$$Z_{J,D} \xleftarrow{\mathfrak{c}} Z_{J,J',D} \xrightarrow{\mathfrak{d}} Z_{J',D}$$

where

$$\begin{aligned} Z_{J,J',D} &= \{(P, gU_Q); P \in \mathcal{P}_J, Q = Q_{J',P}, gU_Q \in D/U_Q\}, \\ \mathfrak{c}(P, gU_Q) &= (P, gU_P), \mathfrak{d}(P, gU_Q) = (Q, gU_Q). \end{aligned}$$

Define functors

$$\mathfrak{f}_{J,J'} : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(Z_{J',D}), \mathfrak{e}_{J,J'} : \mathcal{D}(Z_{J',D}) \rightarrow \mathcal{D}(Z_{J,D})$$

by $\mathfrak{f}_{J,J'} A = \mathfrak{d}_! \mathfrak{c}^* A$, $\mathfrak{e}_{J,J'} A' = \mathfrak{c}_! \mathfrak{d}^* A'$. Now $\mathfrak{d}$ is proper and $\mathfrak{c}$ is an affine space bundle with fibres of dimension $a = \dim \mathcal{P}_J - \dim \mathcal{P}_{J'}$. Hence $\mathfrak{f}_{J,J'}$ commutes with Verdier duality up to a shift and a twist:

$$(a) \quad \mathfrak{D} \mathfrak{f}_{J,J'} = \mathfrak{f}_{J,J'} [[a]] \mathfrak{D} : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(Z_{J',D}).$$

For $J \subset J' \subset J'' \subset \mathbf{I}$ we have (see [L10, 6.2]):

$$(b) \quad \mathfrak{f}_{J,J''} = \mathfrak{f}_{J',J''} \mathfrak{f}_{J,J'}, \quad \mathfrak{e}_{J,J''} = \mathfrak{e}_{J,J'} \mathfrak{e}_{J',J''}.$$

Clearly, $\mathfrak{f}_{J,J} = 1$, $\mathfrak{e}_{J,J} = 1$.

For $J \subset J' \subset \mathbf{I}$ we show that the convolution bifunctor

$$\mathcal{D}(Z_{J,D}) \times \mathcal{D}(Z_{\epsilon(J),D'}) \rightarrow \mathcal{D}(Z_{J,D'D}), \quad A, B \mapsto A * B,$$

(see 32.5) is compatible with the functors $\mathfrak{e}_{J,J'}$ in the following sense: for $A \in \mathcal{D}(Z_{J,D})$, $A' \in \mathcal{D}(Z_{\epsilon(J),D'})$, we have

$$(c) \quad (\mathfrak{e}_{J,J'} A) * (\mathfrak{e}_{J,J'} A') = \mathfrak{e}_{J,J'} (A * A').$$

We have a commutative diagram

$$\begin{array}{ccccc} Z_{J',D} \times Z_{\epsilon(J'),D'} & \xleftarrow{b'_1} & Z'_0 & \xrightarrow{b'_2} & Z_{J',D'D} \\ \mathfrak{d} \times \mathfrak{d}' \uparrow & & f_1 \uparrow & & \mathfrak{d}'' \uparrow \\ Z_{J,J',D} \times Z_{\epsilon(J),\epsilon(J'),D'} & \xleftarrow{f_2} & \underline{Z} & \xrightarrow{f_3} & Z_{J,J',D'D} \\ \mathfrak{c} \times \mathfrak{c}' \downarrow & & f_4 \downarrow & & \mathfrak{c}'' \downarrow \\ Z_{J,D} \times Z_{\epsilon(J),D'} & \xleftarrow{b_1} & Z_0 & \xrightarrow{b_2} & Z_{J,D'D} \end{array}$$

Here $\mathfrak{c}, \mathfrak{d}$ are as above,

$Z_{\epsilon(J'),D'} \xleftarrow{\mathfrak{c}'} Z_{\epsilon(J),\epsilon(J'),D'} \xrightarrow{\mathfrak{d}'} Z_{\epsilon(J'),D'}$, $Z_{J,D'D} \xleftarrow{\mathfrak{c}''} Z_{J,J',D'D} \xrightarrow{\mathfrak{d}''} Z_{J',D'D}$ are the analogous maps when (J, J', D) is replaced by $(\epsilon(J), \epsilon(J'), D')$ or by $(J, J', D'D)$, b_1, b_2 are as in 32.5, b'_1, b'_2 are the analogous maps with J replaced by J' ,

$$\begin{aligned} \underline{Z} &= \{((X, P, gU_P, g'U_{P'}); X \in \mathcal{P}_J, P = Q_{J',X}, gU_P \in D/U_P, \\ &\quad P' = gPg^{-1}, g'U_{P'} \in D'/U_{P'}\}, \end{aligned}$$

$$\begin{aligned}
f_1(X, gU_P, g'U_{P'}) &= (P, gPg^{-1}, g'gPg^{-1}g'^{-1}, gU_P, g'U_{gPg^{-1}}), \\
f_2(X, gU_P, g'U_{P'}) &= ((X, gU_P), (gXg^{-1}, g'U_{gPg^{-1}})), \\
f_3(X, gU_P, g'U_{P'}) &= (X, g'gU_P), \\
f_4(X, gU_P, g'U_{P'}) &= (X, gXg^{-1}, g'gXg^{-1}g'^{-1}, gU_X, g'U_{gXg^{-1}}).
\end{aligned}$$

It is enough to show that

$$b_{2!}b_1^*(\mathfrak{c} \times \mathfrak{c}')_!(\mathfrak{d} \times \mathfrak{d}')^*(A \boxtimes A') = \mathfrak{c}'_!\mathfrak{d}''^*b'_{2!}b_1^*(A \boxtimes A').$$

Since the upper right and lower left squares are cartesian we have

$$b_1^*(\mathfrak{c} \times \mathfrak{c}')_! = f_{4!}f_2^*, \mathfrak{d}''^*b'_{2!} = f_{3!}f_1^*.$$

Hence it is enough to show that

$$b_{2!}f_{4!}f_2^*(\mathfrak{d} \times \mathfrak{d}')^*(A \boxtimes A') = \mathfrak{c}'_!f_{3!}f_1^*b_1^*(A \boxtimes A').$$

This follows from the commutativity of the diagram above: we have $b_{2!}f_{4!} = \mathfrak{c}'_!f_{3!}$ and $f_2^*(\mathfrak{d} \times \mathfrak{d}')^* = f_1^*b_1^*$. This proves (c).

If $A \in \mathcal{D}(Z_{J,D})$, $B \in \mathcal{D}(Z_{\epsilon(J),D'})$, $C \in \mathcal{D}(Z_{\epsilon'\epsilon(J),D''})$ then, from definitions, we have the associativity property:

$$(d) \quad (A * B) * C = A * (B * C).$$

Consider the isomorphism $\partial : Z_{J,D} \xrightarrow{\sim} Z_{\epsilon(J),D^{-1}}$ as in 28.19. Note that the composition $Z_{J,D} \xrightarrow{\partial} Z_{\epsilon(J),D^{-1}} \xrightarrow{\partial} Z_{J,D}$ is the identity map. From definitions, we have for A, B as above:

$$(e) \quad \partial_!(A * B) = (\partial_!B) * (\partial_!A).$$

For $J \subset \mathbf{I}$ we define a functor $\tau : \mathcal{D}(Z_{J,G^0}) \rightarrow \mathcal{D}(\text{point})$ by $\tau(C) = p'_!i^*(C)$ where $i : \mathcal{P}_J \rightarrow Z_{J,G^0}$ is the imbedding $P \mapsto (P, U_P)$ and $p' : \mathcal{P}_J \rightarrow \text{point}$ is the obvious map. We define a bifunctor $(:) : \mathcal{D}(Z_{J,D}) \times \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(\text{point})$ by $(A : B) = p_!(A \otimes B)$ where $p : Z_{J,D} \rightarrow \text{point}$ is the obvious map. As in 32.23(b) we have

$$(f) \quad (A : B) = \tau(A * (\partial_!B)).$$

From definitions we have

$$(g) \quad (A : B) = (\partial_!A : \partial_!B).$$

If $A \in \mathcal{D}(Z_{J,D})$, $B \in \mathcal{D}(Z_{\epsilon(J),D'})$, $C \in \mathcal{D}(Z_{\epsilon'\epsilon(J),D^{-1}D'^{-1}})$ then from (d) we have $\tau((A * B) * C) = \tau(A * (B * C))$. Using (f) we rewrite this as $(A * B : \partial_!C) = (A : \partial_!(B * C))$ or, using (g), as:

$$(h) \quad (A * B : \partial_!C) = (\partial_!A : B * C).$$

For $J \subset J'$, $\mathfrak{c}, \mathfrak{d}$ as above and $A \in \mathcal{D}(Z_{J,D})$, $B \in \mathcal{D}(Z_{J',D})$ we show:

$$(i) \quad (B : f_{J,J'}A) = (e_{J,J'}B : A) \in \mathcal{D}(\text{point}).$$

Let p_1, p_2, p_3 be the obvious maps from $Z_{J',D}, Z_{J,D}, Z_{J,J',D}$ to the point. We must show that $p_{1!}(B \otimes \mathfrak{d}_!\mathfrak{c}^*A) = p_{2!}(\mathfrak{c}_!\mathfrak{d}^*B \otimes A)$. Since $p_{1!}\mathfrak{d}_! = p_{3!} = p_{2!}\mathfrak{c}_!$, both sides are equal to $p_{3!}(\mathfrak{d}^*B \otimes \mathfrak{c}^*A)$. This proves (i).

36.5. For $J \subset J' \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, $w' \in {}^{\epsilon(J')}\mathbf{W}$ we have a diagram

$${}^w Z_{J,D} \xleftarrow{\mathfrak{c}_{w,w'}} {}^{w,w'} Z_{J,J',D} \xrightarrow{\mathfrak{d}_{w,w'}} {}^{w'} Z_{J',D}$$

where

$${}^{w,w'} Z_{J,J',D} = \{(P, gU_Q) \in Z_{J,J',D}; (P, gU_P) \in {}^w Z_{J,D}, (Q, gU_Q) \in {}^{w'} Z_{J',D}\}$$

and $\mathfrak{c}_{w,w'}$, $\mathfrak{d}_{w,w'}$ are the restrictions of $\mathfrak{c}$, $\mathfrak{d}$ in 36.4. Consider the functor

$$\mathfrak{f}_{J,w,J',w'} : \mathcal{D}({}^w Z_{J,D}) \rightarrow \mathcal{D}({}^{w'} Z_{J',D}), A \mapsto \mathfrak{d}_{w,w'}! \mathfrak{c}_{w,w'}^* A.$$

From definitions we see that, for $A \in \mathcal{D}({}^w Z_{J,D})$, we have

$$(a) \quad \mathfrak{f}_{J,w,J',w'}(A) = i_{J',w'}^* \mathfrak{f}_{J,J'}(i_{J,w}! A).$$

36.6. Let $J \subset \mathbf{I}$ and let $w \in {}^{\epsilon(J)}\mathbf{W}$. We have a sequence of affine space bundles

$$(a) \quad {}^w Z_{J,D} \xrightarrow{\vartheta_{J,w}} {}^w Z_{J_1,D} \xrightarrow{\vartheta_{J_1,w}} {}^w Z_{J_2,D} \xrightarrow{\vartheta_{J_2,w}} \dots$$

where $\vartheta_{J,w}$ is as in 36.1, $\vartheta_{J_1,w}$ is the analogous map with J replaced by J_1 (we set $(J_1)_1 = J_2$), $\vartheta_{J_2,w}$ is the analogous map with J replaced by J_2 (we set $(J_2)_1 = J_3$), etc. We have $J \supset J_1 \supset J_2 \supset \dots$. Let $J_\infty = J_r$ for large r . Since $\vartheta_{J_r,w} = 1$ for large r , the composition of the maps (a) is a well defined map $\vartheta : {}^w Z_{J,D} \rightarrow {}^w Z_{J_\infty,D}$. For any $(P, gU_P) \in {}^w Z_{J,D}$ we have

$$\vartheta_{J,w}(P, gU_P) = (P_1, gU_{P_1}), \vartheta_{J_1,w}(P_1, gU_{P_1}) = (P_2, gU_{P_2}), \dots$$

where $P \supset P_1 \supset P_2 \supset \dots$. Let $P_\infty = P_r$ for large r . We show:

(b) *The map $\alpha : {}^{w,w} Z_{J_\infty,J,D} \rightarrow {}^w Z_{J,D}$, $(R, gU_P) \mapsto (P, gU_P)$, is an isomorphism.* ■

We show only that α is a bijection. Let $(P, gU_P) \in {}^w Z_{J,D}$. By [L10, 4.14(b)] there exists $(B, h) \in \mathcal{B} \times gU_P$ such that $B \subset P$ and $\text{pos}(hBh^{-1}, B) = w$. We have also $\text{pos}(gBg^{-1}, B) = w$. Let $R = Q_{J_\infty, B}$. By [L10, 4.14(a)] we have $(R, gU_R) \in {}^w Z_{J_\infty,D}$. We have $(R, gU_P) \in {}^{w,w} Z_{J_\infty,J,D}$ and $\alpha(R, gU_P) = (P, gU_P)$. Thus α is surjective.

We show that α is injective. Let $(R, gU_P) \in {}^{w,w} Z_{J_\infty,J,D}$. Define $P_1, P_2, \dots, P_\infty$ in terms of (P, gU_P) as above. It is enough to show that $R = P_\infty$. Let $P' = gPg^{-1}$, $R' = gRg^{-1}$. Since $(R, gU_R) \in {}^w Z_{J_\infty,D}$ and $\epsilon^{-1}(\text{Ad}(w)J_\infty) = J_\infty$, we see that R, R' have a common Levi and $\text{pos}(R', R) = w$. Hence if B' is any Borel of R' , there exists a Borel B of R such that $\text{pos}(B', B) = w$. If $w_0 = \text{pos}(P', P)$, we have $w = w_0 x$ where $x \in \mathbf{W}_J$ (see [L10, 2.1(b)]) and $l(w_0) + l(x) = l(w)$. Hence we can find a Borel B_1 of G^0 such that $\text{pos}(B', B_1) = w_0$, $\text{pos}(B_1, B) = x$. Since $B \subset R \subset P$ we have $B_1 \subset P$. Since $\text{pos}(B', B_1) = \text{pos}(P', P) = w_0$, we have $B' \subset (P' \cap P)U_{P'}$ (see [L10, 2.7]). Since this holds for any Borel B' of

R' and R' is the union of its Borels, it follows that $R' \subset (P' \cap P)U_{P'}$. Hence $R = g^{-1}R'g \subset (g^{-1}Pg \cap P)U_P = P_1$. Hence $(R, gU_{P_1}) \in {}^{w,w}Z_{J_\infty, J_1, D}$ is well defined. Repeating the previous argument for (R, gU_{P_1}) instead of (R, gU_P) we see that $R \subset P_2$. Continuing in this way we obtain $R \subset P_r$ for any $r \geq 0$. In particular $R \subset P_\infty$. Since R, P_∞ are parabolics of the same type, we see that $R = P_\infty$. This proves (b).

Next we show:

(c) If $y \in {}^{\epsilon(J)}\mathbf{W}$, $y \neq w$, then ${}^{w,y}Z_{J_\infty, J, D} = \emptyset$.

Assume that $(R, gU_P) \in {}^{w,y}Z_{J_\infty, J, D}$. We have $(R, gU_R) \in {}^wZ_{J_\infty, D}$ and by [L10, 4.14(b)] there exists $(B, h) \in \mathcal{B} \times gU_R$ such that $B \subset R$ and $\text{pos}(hBh^{-1}, B) = w$. We have also $\text{pos}(gBg^{-1}, B) = w$. Since $B \subset P$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, we see, using [L10, 4.14(a)], that $(P, gU_P) \in {}^wZ_{J, D}$. Since $(P, gU_P) \in {}^yZ_{J, D}$ and ${}^yZ_{J, D}, {}^wZ_{J, D}$ are disjoint for $y \neq w$, we have a contradiction. This proves (c).

Let $A \in {}^w\hat{Z}_{J, D}$. From definitions we have $A = \underline{\vartheta}^\star(A')$ with $A' \in {}^w\hat{Z}_{J_\infty, D}$. We show:

(d) $\mathfrak{f}_{J_\infty, w, J, w}(A') \cong A[-\delta]$ for some integer δ .

(e) $\mathfrak{f}_{J_\infty, w, J, y}(A') = 0$ for any $y \in {}^{\epsilon(J)}\mathbf{W}$, $y \neq w$.

Now (e) follows immediately from (c). To prove (d) we recall that $\mathfrak{f}_{J_\infty, w, J, w}(A') = \alpha_! \tilde{a}^*(A')$ where $\tilde{a} : {}^{w,w}Z_{J_\infty, J, D} \rightarrow {}^wZ_{J_\infty, D}$ is given by $(R, gU_P) \mapsto (R, gU_R)$. Define $e : {}^wZ_{J, D} \rightarrow {}^{w,w}Z_{J_\infty, J, D}$ by $e(P, gU_P) = (P_\infty, gU_P)$ with P_∞ as above. Clearly, $\tilde{a}e = \underline{\vartheta}$, $\alpha e = 1$. Since α is an isomorphism, we see that

$$\alpha_! \tilde{a}^*(A') = e^* \tilde{a}^*(A') = \underline{\vartheta}^*(A') = \underline{\vartheta}^\star(A')[-\delta] = A[-\delta]$$

where δ is the dimension of any fibre of $\underline{\vartheta}$. This proves (d).

36.7. Let $L \subset J \subset \mathbf{I}$ and let $w \in {}^{\epsilon(J)}\mathbf{W}$, $y \in {}^{\epsilon(L)}\mathbf{W}$. Assume that $\epsilon^{-1}(\text{Ad}(w)J) = J$, $\epsilon^{-1}(\text{Ad}(y)L) = L$. From definitions we have

$$\begin{aligned} {}^{y,w}Z_{L, J, D} = \{ & (R, gU_P); R \in \mathcal{P}_L, P = Q_{J, R}, gU_P \in D/U_P, \\ & \text{pos}(gRg^{-1}, R) = y, \text{pos}(gPg^{-1}, P) = w \}. \end{aligned}$$

We show:

(a) ${}^{y,w}Z_{L, J, D} = \emptyset$ unless $y \in \mathbf{W}_{\epsilon(J)}w = w\mathbf{W}_J$ that is, $w = \min(\mathbf{W}_{\epsilon(J)}y\mathbf{W}_J)$.

Assume that $(R, gU_P) \in {}^{y,w}Z_{L, J, D}$. Let B be a Borel of gRg^{-1} . Then $B \subset gPg^{-1}$. In our case gPg^{-1}, P have a common Levi; hence there exists a Borel B' of P such that $\text{pos}(B, B') = \text{pos}(gPg^{-1}, P) = w$. Similarly, since gRg^{-1}, R have a common Levi, there exists a Borel B'' of R such that $\text{pos}(B, B'') = \text{pos}(gRg^{-1}, R) = y$. Let $z = \text{pos}(B', B'')$. Since $B' \subset P, B'' \subset P$, we have $z \in \mathbf{W}_J$. Since $w \in \mathbf{W}^J$ we have $l(wz) = l(w) + l(z)$ hence from $\text{pos}(B, B') = w, \text{pos}(B', B'') = z$ we deduce $\text{pos}(B, B'') = wz$. Thus $y = wz$ and (a) follows.

Now for $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$, $w' \in {}^{\epsilon'(J)}\mathbf{W}$, $y \in {}^{\epsilon'\epsilon(J)}\mathbf{W}$, the diagram

$Z_{J,D} \times Z_{\epsilon(J),D'} \xleftarrow{b_1} Z_0 \xrightarrow{b_2} Z_{J,D'D}$
(see 32.5) restricts to a diagram

$${}^w Z_{J,D} \times {}^{w'} Z_{\epsilon(J),D'} \xleftarrow{b_{w,w',1}} {}^{w,w',y} Z_0 \xrightarrow{b_{w,w',2}} {}^y Z_{J,D'D}$$

where ${}^{w,w',y} Z_0 = b_1^{-1}({}^w Z_{J,D} \times {}^{w'} Z_{\epsilon(J),D'}) \cap b_2^{-1}({}^y Z_{J,D'D})$. Define a bifunctor $\mathcal{D}({}^w Z_{J,D}) \times \mathcal{D}({}^{w'} Z_{\epsilon(J),D'}) \rightarrow \mathcal{D}({}^y Z_{J,D'D})$ by

$$A, B \mapsto A *_y B = b_{w,w',2}! b_{w,w',1}^*(A \boxtimes B).$$

Let $K \in \mathcal{D}({}^w Z_{J,D})$, $K' \in \mathcal{D}({}^{w'} Z_{\epsilon(J),D'})$. From definitions we see that

$$(b) \quad i_{J,y}^*((i_{J,w}! K) * (i_{\epsilon(J),w'}! K')) = K *_y K'.$$

36.8. In the remainder of this section we assume that $\mathbf{k}$ is an algebraic closure of a finite field. Let $J \subset \mathbf{I}$. As in 31.2 let $\mathcal{A} = \mathbf{Z}[v, v^{-1}]$, v an indeterminate. Let $\mathfrak{K}(Z_{J,D})$ (resp. $\mathfrak{K}'(Z_{J,D})$) be the free $\mathcal{A}$ -module with basis given by the elements in $\mathcal{I}_{J,D}$ (resp. in $\sqcup_{w \in {}^\epsilon(J)} \mathbf{W} \mathcal{I}_{J,w,D}$). We have $\mathfrak{K}'(Z_{J,D}) = \oplus_{w \in {}^\epsilon(J)} \mathbf{W} {}^w \mathfrak{K}'(Z_{J,D})$ where ${}^w \mathfrak{K}'(Z_{J,D})$ is the $\mathcal{A}$ -submodule of $\mathfrak{K}'(Z_{J,D})$ generated by $\mathcal{I}_{J,w,D}$. For $w \in {}^\epsilon(J) \mathbf{W}$,

$$i_{J,w}^* : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}({}^w Z_{J,D}), \quad i_{J,w}^* : \mathcal{D}({}^w Z_{J,D}) \rightarrow \mathcal{D}(Z_{J,D})$$

restrict to functors

$$\mathcal{D}^{cs}(Z_{J,D}) \rightarrow \mathcal{D}^{cs}({}^w Z_{J,D}), \quad \mathcal{D}^{cs}({}^w Z_{J,D}) \rightarrow \mathcal{D}^{cs}(Z_{J,D}),$$

see 36.3(a), 36.3(f). These functors, or rather their mixed analogues, are compatible with the triangular structures hence they induces $\mathcal{A}$ -linear maps

$$\phi_{J,w} : \mathfrak{K}(Z_{J,D}) \rightarrow {}^w \mathfrak{K}'(Z_{J,D}), \quad \phi'_{J,w} : {}^w \mathfrak{K}'(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D}).$$

Clearly, $\phi_{J,w} \phi'_{J,w} = 1 : {}^w \mathfrak{K}'(Z_{J,D}) \rightarrow {}^w \mathfrak{K}'(Z_{J,D})$. For $A \in \mathcal{I}_{J,D}$ regarded as a pure complex of weight 0, we have

$$\phi_{J,w}(A) = \sum_{A_1 \in \mathcal{I}_{J,w,D}} \sum_{j,h} (-1)^j v^h (\text{multiplicity of } A_1 \text{ in } {}^p H^j(i_{J,w}^* A)_h) A_1$$

(the subscript h means "the subquotient of pure weight h "). Define an $\mathcal{A}$ -linear map $\phi : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}'(Z_{J,D})$ by $\phi = \oplus_{w \in {}^\epsilon(J)} \mathbf{W} \phi_{J,w}$.

From 36.3(b),(d) we see that the matrix of ϕ is square and upper triangular (with 1 on diagonal) with respect to a suitable order on $\mathcal{I}_{J,D}$. In particular,

(a) $\phi : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}'(Z_{J,D})$ is an isomorphism.

Note that the inverse isomorphism restricted to ${}^w \mathfrak{K}'(Z_{J,D})$ is just $\phi'_{J,w}$. Clearly,

$$(b) \quad \mathfrak{K}(Z_{J,D}) = \oplus_{w \in {}^\epsilon(J)} \mathbf{W} {}^w \mathfrak{K}(Z_{J,D}),$$

where ${}^w \mathfrak{K}(Z_{J,D}) = \phi^{-1}({}^w \mathfrak{K}'(Z_{J,D}))$ for $w \in {}^\epsilon(J) \mathbf{W}$.

We have a partition $\mathcal{I}_{J,D} = \sqcup_{\mathfrak{k} \in \mathbf{W} \backslash \underline{\mathfrak{s}}(\mathbf{T})} \mathcal{I}_{J,D}^{\mathfrak{k}}$ where $\mathbf{W} \backslash \underline{\mathfrak{s}}(\mathbf{T})$ is the set of $\mathbf{W}$ -orbits on $\underline{\mathfrak{s}}(\mathbf{T})$ and $\mathcal{I}_{J,D}^{\mathfrak{k}}$ consists of those $A \in \mathcal{I}_{J,D}$ such that $A \in \hat{Z}_{J,D}^{\mathcal{L}}$ for some $\mathcal{L} \in \mathfrak{s}(\mathbf{T})$ whose isomorphism class is in $\mathfrak{k}$. We have

$$(c) \mathfrak{K}(Z_{J,D}) = \bigoplus_{\mathfrak{k} \in \mathbf{W} \setminus \underline{\mathfrak{s}}(\mathbf{T})} \mathfrak{K}^{\mathfrak{k}}(Z_{J,D})$$

where $\mathfrak{K}^{\mathfrak{k}}(Z_{J,D})$ is the $\mathcal{A}$ -submodule of $\mathfrak{K}(Z_{J,D})$ generated by $\mathcal{I}_{J,D}^{\mathfrak{k}}$. From definitions we see that each ${}^w\mathfrak{K}'(Z_{J,D})$ and each ${}^w\mathfrak{K}(Z_{J,D})$ has a natural direct sum decomposition indexed by $\mathbf{W} \setminus \underline{\mathfrak{s}}(\mathbf{T})$ analogous to (c). It follows that the decompositions (b),(c) are compatible in the sense that

$$(d) \mathfrak{K}(Z_{J,D}) = \bigoplus_{w, \mathfrak{k}} ({}^w\mathfrak{K}(Z_{J,D}) \cap \mathfrak{K}^{\mathfrak{k}}(Z_{J,D})).$$

36.9. Let $J \subset J' \subset \mathbf{I}$. As in [L10, 6.4, 6.7(b)] we see that $\mathfrak{f}_{J,J'}, \mathfrak{e}_{J,J'}$ restrict to functors $\mathfrak{f}_{J,J'} : \mathcal{D}^{cs}(Z_{J,D}) \rightarrow \mathcal{D}^{cs}(Z_{J',D})$, $\mathfrak{e}_{J,J'} : \mathcal{D}^{cs}(Z_{J',D}) \rightarrow \mathcal{D}^{cs}(Z_{J,D})$, denoted again by $\mathfrak{f}_{J,J'}, \mathfrak{e}_{J,J'}$. These functors or, rather, their mixed analogues, are compatible with the triangulated structures hence induce $\mathcal{A}$ -linear maps $\mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J',D})$, $\mathfrak{K}(Z_{J',D}) \rightarrow \mathfrak{K}(Z_{J,D})$ denoted again by $\mathfrak{f}_{J,J'}, \mathfrak{e}_{J,J'}$; the identities 36.4(b) continue to hold for these linear maps. Moreover, Verdier duality induces a group homomorphism $\mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ (denoted again by $\mathfrak{D}$) and from 36.4(a) we deduce

$$(a) \quad \mathfrak{D}\mathfrak{f}_{J,J'} = v^{-2a}\mathfrak{f}_{J,J'}\mathfrak{D} : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J',D})$$

with a as in 36.4(a). Now for any $A \in \mathcal{I}_{J,D}$ we have

$$(b) \quad \mathfrak{D}(v^m A) = v^{-m} A^*$$

where $A^* \in \mathcal{I}_{J,D}$ is isomorphic to the Verdier dual of A . If $J \subset J'$, $\mathfrak{c}, \mathfrak{d}, a$ are as in 36.4, then for $A \in \mathcal{I}_{J,D}, B \in \mathcal{I}_{J',D}$ (regarded as pure perverse sheaves of weight 0) we have

$$\mathfrak{f}_{J,J'} A = \sum_{A_1 \in \mathcal{I}_{J',D}} \sum_{j,h} (-1)^j v^h (\text{multiplicity of } A_1 \text{ in } {}^p H^j(\mathfrak{d}! \mathfrak{c}^* A)_h) A_1,$$

$$\mathfrak{e}_{J,J'} B = \sum_{B_1 \in \mathcal{I}_{J,D}} \sum_{j,h} (-1)^j v^h (\text{multiplicity of } B_1 \text{ in } {}^p H^j(\mathfrak{c}! \mathfrak{d}^* B)_h) B_1$$

(the subscript h means "the subquotient of pure weight h "). Since $\mathfrak{c}$ is an affine space bundle with fibres of dimension a , $\mathfrak{c}^* A[a]$ is a pure perverse sheaf of weight a . By [De, 6.2.6] applied to the proper morphism $\mathfrak{d}$, $\mathfrak{d}! \mathfrak{c}^* A[a]$ is a pure complex of weight a . From [BBD, 5.4.1], we see that ${}^p H^j(\mathfrak{d}! \mathfrak{c}^* A) = {}^p H^{j-a}(\mathfrak{d}! \mathfrak{c}^* A[a])$ is pure of weight $(j-a) + a = j$. Hence, writing $\mathfrak{f}_{J,J'} A = \sum_{A_1 \in \mathcal{I}_{J',D}} x_{A,A_1} A_1$, $x_{A,A_1} \in \mathcal{A}$ we have

$$x_{A,A_1} = \sum_j (-v)^j (\text{multiplicity of } A_1 \text{ in } {}^p H^{j-a}(\mathfrak{d}! \mathfrak{c}^* A[a])).$$

Let $\xi \mapsto \bar{\xi}$ be the group homomorphism $\mathcal{A} \rightarrow \mathcal{A}$ given by $v^m \mapsto v^{-m}$ for $m \in \mathbf{Z}$. Since $\mathfrak{c}^* A[a]$ is a pure perverse sheaf we have (using the relative hard Lefschetz

theorem [BBD, 5.4.10]):

$$\begin{aligned}
\bar{x}_{A,A_1} &= \sum_j (-v)^{-j} (\text{multiplicity of } A_1 \text{ in } {}^p H^{j-a}(\mathfrak{d}_! \mathfrak{c}^* A[a])) \\
&= \sum_j (-v)^{-j} (\text{multiplicity of } A_1 \text{ in } {}^p H^{-j+a}(\mathfrak{d}_! \mathfrak{c}^* A[a])) \\
&= \sum_{j'} (-v)^{j'-2a} (\text{multiplicity of } A_1 \text{ in } {}^p H^{j'-a}(\mathfrak{d}_! \mathfrak{c}^* A[a])) = v^{-2a} x_{A,A_1}.
\end{aligned}$$

Define a group homomorphism $\beta_J : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ by $\beta_J(v^m A) = v^{-m} A$ for $A \in \mathcal{I}_{J,D}, m \in \mathbf{Z}$. We see that $\beta_{J'}(\mathfrak{f}_{J,J'} A) = v^{-2a} \mathfrak{f}_{J,J'} A$ for any $A \in \mathcal{I}_{J,D}, m \in \mathbf{Z}$. Hence for any $\xi \in \mathfrak{K}(Z_{J,D})$ we have

$$(c) \quad \beta_{J'}(\mathfrak{f}_{J,J'} \xi) = v^{-2a} \mathfrak{f}_{J,J'}(\beta_J(\xi)).$$

Define $\tilde{\mathfrak{D}} : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ by $\tilde{\mathfrak{D}} A = \mathfrak{D} \beta_J(A) = \beta_J \mathfrak{D}(A)$. Clearly, $\tilde{\mathfrak{D}}$ is $\mathcal{A}$ -linear. Note that

(d) *the maps $\mathfrak{f}_{J,J'}, \mathfrak{e}_{J,J'}, \beta_J$ are compatible with the decompositions of type 36.8(c) while $\mathfrak{D}, \tilde{\mathfrak{D}}$ map the summand corresponding to $\mathfrak{k}$ in 36.8(c) onto the summand corresponding to $\tilde{\mathfrak{k}}$. (Here $\tilde{\mathfrak{k}}$ is the image of $\mathfrak{k}$ under $\mathcal{L} \mapsto \tilde{\mathcal{L}}$, see 28.18.)*

From 32.21 we see that convolution (see 32.5) restricts to a bifunctor

$$\mathcal{D}^{cs}(Z_{J,D}) \times \mathcal{D}^{cs}(Z_{\epsilon(J),D'}) \rightarrow \mathcal{D}^{cs}(Z_{J,D'D}).$$

This induces an $\mathcal{A}$ -bilinear pairing $\mathfrak{K}(Z_{J,D}) \times \mathfrak{K}(Z_{\epsilon(J),D'}) \rightarrow \mathfrak{K}(Z_{J,D'D})$. It sends $A \in \mathcal{I}_{J,D}, B \in \mathcal{I}_{\epsilon(J),D'}$ (regarded as pure perverse sheaves of weight 0) to

$$\sum_{A_1 \in \mathcal{I}_{J,D'D}} \sum_{j,h} (-1)^j v^h (\text{multiplicity of } A_1 \text{ in } {}^p H^j(A * B)_h) A_1.$$

This bilinear pairing is denoted again by $A, B \mapsto A * B$. If $\mathfrak{k}, \mathfrak{k}' \in \mathbf{W} \backslash \underline{\mathfrak{s}}(\mathbf{T})$ and $\xi \in \mathfrak{K}^{\mathfrak{k}}(Z_{J,D}), \xi'^{\mathfrak{k}'} \in \mathfrak{K}(Z_{\epsilon(J),D'})$, then $\xi * \xi' \in \mathfrak{K}^{\mathfrak{k}}(Z_{J,D'D})$ if $\mathfrak{k}' = \underline{D}(\mathfrak{k})$ and $\xi * \xi' = 0$ if $\mathfrak{k}' \neq \underline{D}(\mathfrak{k})$ (see 32.6(a),(b)).

From 28.19 we see that $\partial_! : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(Z_{\epsilon(J),D-1})$ restricts to an equivalence of categories $\mathcal{D}^{cs}(Z_{J,D}) \rightarrow \mathcal{D}^{cs}(Z_{\epsilon(J),D-1})$; this induces an $\mathcal{A}$ -linear isomorphism $\mathfrak{K}(Z_{J,D}) \xrightarrow{\sim} \mathfrak{K}(Z_{\epsilon(J),D-1})$ denoted again by ∂ . Now the composition $\mathfrak{K}(Z_{J,D}) \xrightarrow{\partial} \mathfrak{K}(Z_{\epsilon(J),D-1}) \xrightarrow{\partial} \mathfrak{K}(Z_{J,D})$ is the identity map.

In the setup of 36.5, $\mathfrak{f}_{J,w,J',w'}$ restricts to a functor $\mathfrak{f}_{J,w,J',w'} : \mathcal{D}^{cs}({}^w Z_{J,D}) \rightarrow \mathcal{D}^{cs}({}^{w'} Z_{J',D})$. (We use the analogous statement for $\mathfrak{f}_{J,J'}$, as above, together with 36.5(a), 36.3(a), 36.3(f).) Hence there is an induced $\mathcal{A}$ -linear map ${}^w \mathfrak{K}'(Z_{J,D}) \rightarrow {}^{w'} \mathfrak{K}'(Z_{J',D})$ denoted again by $\mathfrak{f}_{J,w,J',w'}$. Let $\mathfrak{f}'_{J,J'} : \mathfrak{K}'(Z_{J,D}) \rightarrow \mathfrak{K}'(Z_{J',D})$ be the unique $\mathcal{A}$ -linear map such that $pr_{w'} \mathfrak{f}'_{J,J'} pr_w = \mathfrak{f}_{J,w,J',w'}$ for any w, w' as above. (Here pr_w is the projection of $\mathfrak{K}'(Z_{J',D})$ onto the direct summand ${}^w \mathfrak{K}'(Z_{J,D})$.) From 36.5(a) we see that for $x \in {}^w \mathfrak{K}'(Z_{J',D})$ we have

$$(e) \quad \mathfrak{f}'_{J,J'} = \phi \mathfrak{f}_{J,J'} \phi^{-1}.$$

We also have

$$(f) \quad \partial \mathfrak{f}_{J,J'} = \mathfrak{f}_{\epsilon(J),\epsilon(J')} \partial, \partial \mathfrak{e}_{J,J'} = \mathfrak{e}_{\epsilon(J),\epsilon(J')} \partial.$$

Now the functor $\tau : \mathcal{D}^{cs}(Z_{J,G^0}) \rightarrow \mathcal{D}(\text{point})$ (see 36.4) or rather its mixed analogue induces an $\mathcal{A}$ -linear map $\mathfrak{K}(Z_{J,G^0}) \rightarrow \mathcal{A}$; it sends $C \in \mathcal{I}_{J,G^0}$ regarded as a pure complex of weight 0 to

$$\sum_{j,h} (-1)^j v^h \dim(H_c^j(\mathcal{P}_J, i^*C)_h) \in \mathcal{A}$$

where $i : \mathcal{P}_J \rightarrow Z_{J,G^0}$ is $P \mapsto (P, U_P)$ and $H_c^j(\mathcal{P}_J, i^*C)_h$ denotes the part of weight h of the mixed vector space $H_c^j(\mathcal{P}_J, i^*C)$. This linear map is denoted again by τ .

Similarly, the bifunctor $(:) : \mathcal{D}^{cs}(Z_{J,D}) \times \mathcal{D}^{cs}(Z_{J,D}) \rightarrow \mathcal{D}(\text{point})$ (see 36.4) or rather its mixed analogue induces an $\mathcal{A}$ -bilinear pairing $\mathfrak{K}(Z_{J,D}) \times \mathfrak{K}(Z_{J,D}) \rightarrow \mathcal{A}$; it sends $A, B \in \mathcal{I}_{J,D}$ regarded as pure complexes of weight 0 to

$$\sum_{j,h} (-1)^j v^h \dim(H_c^j(Z_{J,D}, A \otimes B)_h) \in \mathcal{A}.$$

(Here $H_c^j(Z_{J,D}, A \otimes B)_h$ denotes the part of weight h of the mixed vector space $H_c^j(Z_{J,D}, A \otimes B)$.) This bilinear pairing is denoted again by $(:)$. From [BBD, 5.1.14] we see that $(A : B) \in \mathbf{Z}[v^{-1}]$ and from [L3, II, 7.4] and its proof we see that the constant term of $(A : B)$ is δ_{A,B^*} . Thus,

$$(g) \quad (A : B) \in \delta_{A,B^*} + v^{-1} \mathbf{Z}[v^{-1}].$$

We show:

(h) *if $\xi \in \mathfrak{K}(Z_{J,D})$, $(\xi : \tilde{\mathfrak{D}}(\xi)) = 0$ then $\xi = 0$.*

Assume that $\xi \neq 0$. We have $\xi = \sum_{A \in \mathcal{I}_{J,D}, m \in \mathbf{Z}} g_{A,m} v^m A$ where $g_{A,m} \in \mathbf{Z}$ is zero for all but finitely many A, m . We can find $e \in \mathbf{Z}$ such that $g_{A,e} \neq 0$ for some $A \in \mathcal{I}_{J,D}$ and $g_{A,m} = 0$ for all $m > e$ and all $A \in \mathcal{I}_{J,D}$. Then

$$(\xi : \tilde{\mathfrak{D}}(\xi)) = \sum_{A_1, A, m_1, m} g_{A_1, m_1} g_{A, m} v^{m_1 + m} (A_1 : A^*).$$

By (g) this equals $\sum_A g_{A,e}^2 v^{2e} + \text{an element in } v^{2e-1} \mathbf{Z}[v^{-1}]$. This is non-zero since $\sum_A g_{A,e}^2 \in \mathbf{Z}_{>0}$. This proves (h).

We show:

(i) *If $\mathfrak{k}, \mathfrak{k}' \in \mathbf{W} \backslash \mathfrak{s}(\mathbf{T})$ and $\mathfrak{k}' \neq \check{\mathfrak{k}}$ then $(\mathfrak{K}^{\mathfrak{k}}(Z_{J,D}) : \mathfrak{K}^{\mathfrak{k}'}(Z_{J,D})) = 0$.*

Let $A \in \hat{Z}_{J,D}^{\mathcal{L}}, B \in \hat{Z}_{J,D}^{\mathcal{L}_1}$ where $\mathcal{L}, \mathcal{L}_1 \in \mathfrak{s}(\mathbf{T})$. It is enough to show that:

if $H_c^j(Z_{J,D}, A \otimes B) \neq 0$ for some j then the isomorphism class of $\mathcal{L}_1$ is in the $\mathbf{W}$ -orbit of the isomorphism class of $\check{\mathcal{L}}$.

From our assumption we have $A * \partial_!(B) \neq 0$. We have $A \dashv \bar{K}_{J,D}^{\bar{s}, \mathcal{L}}$ as in 28.13(v) and similarly $B \dashv \bar{K}_{J,D}^{\bar{s}', \mathcal{L}_1}$. Hence $\partial_! B \dashv \bar{K}_{\epsilon(J), D-1}^{\bar{s}', \mathcal{L}''}$ where $\mathcal{L}'' = (\underline{D}^{-1})^* \check{\mathcal{L}}$, see 28.19.

We then have $\bar{K}_{J,D}^{\bar{s}, \mathcal{L}} * \bar{K}_{\epsilon(J), D-1}^{\bar{s}', \mathcal{L}''} \neq 0$. Using 32.6(a) we see that $\mathcal{L}, \mathcal{L}_1$ have the required property. This proves (i).

We show:

(j) *If $w, w' \in \mathbf{W}$ and $w \neq w'$ then $({}^w \mathfrak{K}(Z_{J,D}) : {}^{w'} \mathfrak{K}(Z_{J,D})) = 0$.*

It is enough to show: if $A \in \mathcal{D}^{cs}({}^w Z_{J,D}), A' \in \mathcal{D}^{cs}({}^{w'} Z_{J,D})$ then $p_!(i_{J,w}! A \otimes i_{J,w'}! A') = 0$ with p as in 36.4. It is enough to show that $(i_{J,w} \times i_{J,w'})! f^*(A \boxtimes A') = 0$

where $f : Z_{J,D} \rightarrow Z_{J,D} \times Z_{J,D}$ is the diagonal map. This follows from the fact that ${}^w Z_{J,D} \cap ({}^{w'} Z_{J,D}) = \emptyset$.

We show:

(k) If $w \in {}^{\epsilon(J)}\mathbf{W}$, $w \neq 1$ then $\tau({}^w \mathfrak{K}(Z_{J,G^0})) = 0$.

Since ${}^w Z_{J,G^0}, {}^1 Z_{J,G^0}$ are disjoint, we have $i_{J,1}^* i_{J,w!} = 0$ hence $i^* i_{J,w!} = 0$ with i as in 36.4; (k) follows.

From the definitions we have

$$(l) \partial({}^w \mathfrak{K}(Z_{J,D})) = {}^{w^{-1}} \mathfrak{K}(Z_{\epsilon(J), D^{-1}})$$

for any $w \in {}^{\epsilon(J)}\mathbf{W}$ such that $\epsilon^{-1}(\text{Ad}(w)J) = J$. (For such w we have $w^{-1} \in {}^J\mathbf{W}$.)

36.10. Let

$$\tilde{\mathfrak{K}}(Z_{J,D}) = \sum_{L; L \subsetneq J} \mathfrak{f}_{L,J}(\mathfrak{K}(Z_{L,D})), \quad \mathfrak{K}_0 = \bigoplus_{w \in {}^{\epsilon(J)}\mathbf{W}; \epsilon^{-1}(\text{Ad}(w)J) \neq J} {}^w \mathfrak{K}(Z_{J,D}).$$

We show:

$$(a) \quad \tilde{\mathfrak{K}}(Z_{J,D}) = \mathfrak{K}_0 \oplus \bigoplus_{\substack{w \in {}^{\epsilon(J)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} (\tilde{\mathfrak{K}}(Z_{J,D}) \cap {}^w \mathfrak{K}(Z_{J,D})).$$

Setting

$$\tilde{\mathfrak{K}}'(Z_{J,D}) = \sum_{L; L \subsetneq J} \mathfrak{f}'_{L,J}(\mathfrak{K}'(Z_{L,D})), \quad \mathfrak{K}'_0 = \bigoplus_{w \in {}^{\epsilon(J)}\mathbf{W}; \epsilon^{-1}(\text{Ad}(w)J) \neq J} {}^w \mathfrak{K}'(Z_{J,D})$$

and using 36.9(e), we see that it is enough to show:

$$(b) \quad \tilde{\mathfrak{K}}'(Z_{J,D}) = \mathfrak{K}'_0 \oplus \bigoplus_{\substack{w \in {}^{\epsilon(J)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} (\tilde{\mathfrak{K}}'(Z_{J,D}) \cap {}^w \mathfrak{K}'(Z_{J,D})).$$

From 36.6(d), 36.6(e) we see that, if $w \in {}^{\epsilon(J)}\mathbf{W}$, $\epsilon^{-1}(\text{Ad}(w)J) \neq J$, then $A = \mathfrak{f}'_{J_\infty, J} A'$ for some $A' \in {}^y \mathfrak{K}'(Z_{L',D})$ and $J_\infty \neq J$. Hence

$$(c) \quad \mathfrak{K}'_0 \subset \tilde{\mathfrak{K}}'(Z_{J,D}).$$

Using again 36.6(d), 36.6(e) and also 36.4(b) (or rather its analogue for $\mathfrak{f}'_{J,J'}$) and (c), we see that

$$\begin{aligned} \tilde{\mathfrak{K}}'(Z_{J,D}) &= \sum_{\substack{L; L \subsetneq J \\ y \in {}^{\epsilon(L)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(y)L) = L}} \mathfrak{f}'_{L,J}({}^y \mathfrak{K}'(Z_{L,D})) \\ &= \mathfrak{K}'_0 + \sum_{\substack{L; L \subsetneq J \\ y \in {}^{\epsilon(L)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(y)L) = L}} \mathfrak{f}'_{L,J}({}^y \mathfrak{K}'(Z_{L,D})). \end{aligned}$$

Thus, (b) holds and (a) holds.

Let

$$\bar{\mathfrak{K}}(Z_{J,D}) = \mathfrak{K}(Z_{J,D}) / \tilde{\mathfrak{K}}(Z_{J,D}).$$

For any $w \in {}^{\epsilon(J)}\mathbf{W}$ such that $\epsilon^{-1}(\text{Ad}(w)J) = J$, let ${}^w \bar{\mathfrak{K}}(Z_{J,D})$ be the image of ${}^w \mathfrak{K}(Z_{J,D})$ under the obvious map $\mathfrak{K}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D})$. From (a) we see that

$$(d) \quad \bar{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\substack{w \in {}^{\epsilon(J)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} {}^w \bar{\mathfrak{K}}(Z_{J,D}).$$

For any $\mathfrak{k} \in \mathbf{W} \backslash \underline{\mathfrak{s}}(\mathbf{T})$ let $\bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$ be the image of $\mathfrak{K}^{\mathfrak{k}}(Z_{J,D})$ under the obvious map $\mathfrak{K}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D})$. From 36.9(d) we see that $\bar{\mathfrak{K}}(Z_{J,D})$ is compatible with the decomposition 36.8(c). It follows that

$$(e) \quad \bar{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\mathfrak{k} \in \mathbf{W} \backslash \underline{\mathfrak{s}}(\mathbf{T})} \bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D}).$$

Using 36.8(d) we see that the decompositions (d),(e) are compatible in the sense that

$$(f) \quad \bar{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{w, \mathfrak{k}} ({}^w \bar{\mathfrak{K}}(Z_{J,D}) \cap \bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})).$$

Using 36.9(a), 36.9(c), we see that

(g) $\mathfrak{D}, \beta_J, \tilde{\mathfrak{D}} : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ map $\tilde{\mathfrak{K}}(Z_{J,D})$ into itself hence induces group homomorphisms $\bar{\mathfrak{K}}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D})$ denoted again by $\mathfrak{D}, \beta_J, \tilde{\mathfrak{D}}$.

36.11. For $J' \subset J \subset \mathbf{I}$ and $x \in \mathfrak{K}(Z_{J',D}), x' \in \mathfrak{K}(Z_{\epsilon(J),D'})$, we show:

$$(a) \quad (\mathfrak{f}_{J',J}x) * x' = \mathfrak{f}_{J',J}(x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x').$$

Using successively 36.4(h), 36.9(f), 36.4(i), 36.4(c), 36.9(f), 36.4(h), 36.4(i), we see that, for any $x'' \in \mathfrak{K}(Z_{J,D'D})$, we have

$$\begin{aligned} ((\mathfrak{f}_{J',J}x) * x' : x'') &= ((\partial \mathfrak{f}_{J',J}x) : x' * \partial x'') \\ &= (\mathfrak{f}_{\epsilon(J'),\epsilon(J)}\partial(x) : x' * (\partial x'')) = (\partial x : \mathfrak{e}_{\epsilon(J'),\epsilon(J)}(x' * (\partial x''))) \\ &= (\partial x : \mathfrak{e}_{\epsilon(J'),\epsilon(J)}(x') * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}(\partial x'')) = (\partial x : \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x' * \partial \mathfrak{e}_{J',J}x'') \\ &= (x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x' : \mathfrak{e}_{J',J}x'') = (\mathfrak{f}_{J',J}(x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x') : x''). \end{aligned}$$

Thus, if $\xi = (\mathfrak{f}_{J',J}x) * x' - \mathfrak{f}_{J',J}(x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x')$ then $(\xi : x'') = 0$ for any $x'' \in \mathfrak{K}(Z_{J,D'D})$. In particular, $(\xi : \tilde{\mathfrak{D}}(\xi)) = 0$. Using 36.9(h) we see that $\xi = 0$. This proves (a).

A similar argument shows that, if $x \in \mathfrak{K}(Z_{J,D}), x' \in \mathfrak{K}(Z_{\epsilon(J'),D'})$, then:

$$(b) \quad x * (\mathfrak{f}_{\epsilon(J'),\epsilon(J)}x') = \mathfrak{f}_{J',J}(\mathfrak{e}_{J',J}x * x').$$

From (a),(b) we see that

$$\tilde{\mathfrak{K}}(Z_{J,D}) * \mathfrak{K}(Z_{\epsilon(J),D'}) \subset \tilde{\mathfrak{K}}(Z_{J,D'D}), \quad \mathfrak{K}(Z_{J,D}) * \tilde{\mathfrak{K}}(Z_{\epsilon(J),D'}) \subset \tilde{\mathfrak{K}}(Z_{J,D'D}).$$

It follows that $\mathfrak{K}(Z_{J,D}) \times \mathfrak{K}(Z_{\epsilon(J),D'}) \rightarrow \mathfrak{K}(Z_{J,D'D}), A, B \mapsto A * B$, induces an $\mathcal{A}$ -bilinear pairing $\bar{\mathfrak{K}}(Z_{J,D}) \times \bar{\mathfrak{K}}(Z_{\epsilon(J),D'}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D'D})$. We denote it again by $A, B \mapsto A * B$. The following result relates this bilinear pairing to the decompositions of type 36.10(d).

For $w \in {}^{\epsilon(J)}\mathbf{W}, w' \in {}^{\epsilon'\epsilon(J)}\mathbf{W}$ such that $\epsilon^{-1}(\text{Ad}(w)J) = J, \epsilon'^{-1}(\text{Ad}(w')\epsilon(J)) = \epsilon(J)$, let $X_{w,w'}$ be the set of all $y \in {}^{\epsilon'\epsilon(J)}\mathbf{W}$ such that $\epsilon^{-1}\epsilon'^{-1}(\text{Ad}(y)J) = J$ and such that for some $Q \in \mathcal{P}_J, Q' \in \mathcal{P}_{\epsilon(J)}, Q'' \in \mathcal{P}_{\epsilon'\epsilon(J)}$ we have

$$\text{pos}(Q', Q) = w, \text{pos}(Q'', Q') = w', \text{pos}(Q'', Q) = y.$$

Then

$$(c) \quad {}^w \bar{\mathfrak{K}}(Z_{J,D}) * {}^{w'} \bar{\mathfrak{K}}(Z_{\epsilon(J),D'}) \subset \bigoplus_{y \in X_{w,w'}} {}^y \bar{\mathfrak{K}}(Z_{J,D'D}).$$

Indeed, using 36.7(a) we see that it is enough to show:

If ${}^{w,w',y}Z_0 \neq \emptyset$ (notation of 36.7) then $y \in X_{w,w'}$.

This is immediate from definitions.

36.12. As a special case of the pairing $A, B \mapsto A * B$ in 36.11 we have an $\mathcal{A}$ -bilinear pairing $\bar{\mathfrak{K}}(Z_{J,G^0}) \times \bar{\mathfrak{K}}(Z_{J,G^0}) \rightarrow \bar{\mathfrak{K}}(Z_{J,G^0})$. This defines an associative algebra structure on $\bar{\mathfrak{K}}(Z_{J,G^0})$ (not necessarily with 1).

Also as special cases of the pairing $A, B \mapsto A * B$ in 36.11 we have $\mathcal{A}$ -bilinear pairings

$$\bar{\mathfrak{K}}(Z_{J,G^0}) \times \bar{\mathfrak{K}}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D}), \bar{\mathfrak{K}}(Z_{J,D}) \times \bar{\mathfrak{K}}(Z_{\epsilon(J),G^0}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D}),$$

which make $\bar{\mathfrak{K}}(Z_{J,D})$ into a (not necessarily unital) $(\bar{\mathfrak{K}}(Z_{J,G^0}), \bar{\mathfrak{K}}(Z_{\epsilon(J),G^0}))$ -bimodule. ■

By 36.10(f) we have

$$\bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0}) = \bigoplus_{\substack{w \in \epsilon(J)\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} ({}^w \bar{\mathfrak{K}}(Z_{J,G^0}) \cap \bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0})).$$

From 36.10(g) we see that $\mathfrak{D}, \beta_J, \tilde{\mathfrak{D}}$ may be regarded as group homomorphisms $\bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0}) \rightarrow \bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0})$.

36.13. For any $\mathcal{A}$ -module V we set $\underline{V} = \mathbf{Q}(v) \otimes_{\mathcal{A}} V$. If V, V' are $\mathcal{A}$ -modules and $f : V \rightarrow V'$ is $\mathcal{A}$ -linear we denote again by f the $\mathbf{Q}(v)$ -linear map $\underline{V} \rightarrow \underline{V}'$ such that $1 \otimes x \mapsto 1 \otimes f(x)$ for $x \in V$.

In particular, the $\mathbf{Q}(v)$ -vector spaces $\underline{\mathfrak{K}}(Z_{J,D}), \tilde{\underline{\mathfrak{K}}}(Z_{J,D})$ and $\bar{\underline{\mathfrak{K}}}(Z_{J,D})$ are defined. Note that

$$\tilde{\underline{\mathfrak{K}}}(Z_{J,D}) = \sum_{L; L \subsetneq J} \mathfrak{f}_{L,J}(\tilde{\underline{\mathfrak{K}}}(Z_{L,D})), \quad \underline{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\mathfrak{k} \in \mathbf{W} \setminus \underline{\mathfrak{g}}(\mathbf{T})} \underline{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D}),$$

see 36.8(c).

The symmetric bilinear form $(:): \mathfrak{K}(Z_{J,D}) \times \mathfrak{K}(Z_{J,D}) \rightarrow \mathcal{A}$ extends uniquely to a symmetric $\mathbf{Q}(v)$ -bilinear form $\underline{\mathfrak{K}}(Z_{J,D}) \times \underline{\mathfrak{K}}(Z_{J,D}) \rightarrow \mathbf{Q}(v)$ denoted again by $(:)$.

A vector subspace E of $\underline{\mathfrak{K}}(Z_{J,D})$ is said to be *homogeneous* if $E = \sum_{\mathfrak{k} \in \mathbf{W} \setminus \underline{\mathfrak{g}}(\mathbf{T})} E^{\mathfrak{k}}$ where $E^{\mathfrak{k}} = E \cap \underline{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$. For such E we set $E^\perp = \{x \in \underline{\mathfrak{K}}(Z_{J,D}); (x : E) = 0\}$. Then $E^\perp$ is homogeneous. We show that if, in addition, E is stable under $\tilde{\mathfrak{D}}$, then:

$$(a) \quad \underline{\mathfrak{K}}(Z_{J,D}) = E \oplus E^\perp.$$

We first show that $E \cap E^\perp = 0$. Assume that $x \in E \cap E^\perp$. Then $\tilde{f}\tilde{\mathfrak{D}}(x) \in E$ and $(x : \tilde{\mathfrak{D}}(x)) = 0$. We can find $\lambda \in \mathcal{A} - \{0\}$ such that $\lambda x \in \mathfrak{K}(Z_{J,D})$. Then $(\lambda x, \tilde{\mathfrak{D}}(\lambda x)) = 0$. By 36.9(h) we have $\lambda x = 0$. Hence $x = 0$ so that $E \cap E^\perp = 0$. It remains to show that $\dim(E^\perp)^{\mathfrak{k}} + \dim E^{\mathfrak{k}} = \dim \underline{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$ for any $\mathfrak{k}$. This is a consequence of the following statement:

(b) *for any $\mathfrak{k}$, the form $(:)$ on the finite dimensional vector space $\underline{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D}) + \tilde{\underline{\mathfrak{K}}}^{\mathfrak{k}}(Z_{J,D})$ is non-singular.*

More generally we show that for any finite dimensional subspace E' of $\underline{\mathfrak{K}}(Z_{J,D})$ which is stable under $\tilde{\mathfrak{D}}$, the form $(:)$ on E' is non-singular. Let $x \in E'$ be such that $(x : x') = 0$ for any $x' \in E'$. In particular we have $(x, \tilde{\mathfrak{D}}(x)) = 0$. As above, this implies that $x = 0$. This proves (b) hence also (a).

We set

$$\underline{\mathfrak{K}}(Z_{J,D})^J = \{\xi \in \underline{\mathfrak{K}}(Z_{J,D}); \mathfrak{e}_{H,J}\xi = 0 \quad \forall H \subsetneq J\}.$$

We show:

$$(c) \quad \tilde{\mathfrak{K}}(Z_{J,D})^\perp = \mathfrak{K}(Z_{J,D})^J.$$

We have

$$\begin{aligned} \tilde{\mathfrak{K}}(Z_{J,D}) &= \{x \in \mathfrak{K}(Z_{J,D}); (x : \mathfrak{f}_{H,J}\mathfrak{K}(Z_{H,D})) = 0 \quad \forall H \subsetneq J\} \\ &= \{\xi \in \mathfrak{K}(Z_{J,D}); (\mathfrak{e}_{H,J}x : \mathfrak{K}(Z_{H,D})) = 0 \quad \forall H \subsetneq J\} \\ &= \{\xi \in \mathfrak{K}(Z_{J,D}); \mathfrak{e}_{H,J}x = 0 \quad \forall H \subsetneq J\}, \end{aligned}$$

as required. (We have used that $\mathfrak{K}(Z_{H,D})^\perp = 0$ which follows from (a), applied to H instead of J .) This proves (c).

We show:

$$(d) \quad \mathfrak{K}(Z_{J,D}) = \tilde{\mathfrak{K}}(Z_{J,D}) \oplus \mathfrak{K}(Z_{J,D})^J.$$

This follows from (a),(c) using the fact that $\tilde{\mathfrak{K}}(Z_{J,D})$ is stable under $\tilde{\mathfrak{D}}$, see 36.10(g).

From (d) we see that the second projection $\mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})^J$ induces an isomorphism $\mathfrak{K}(Z_{J,D})/\tilde{\mathfrak{K}}(Z_{J,D}) \xrightarrow{\sim} \mathfrak{K}(Z_{J,D})^J$ that is, an isomorphism

$$(e) \quad \bar{\mathfrak{K}}(Z_{J,D}) \xrightarrow{\sim} \mathfrak{K}(Z_{J,D})^J.$$

36.14. The $\mathcal{A}$ -algebra structure on $\mathfrak{K}(Z_{J,G^0})$ (resp. $\bar{\mathfrak{K}}(Z_{J,G^0})$) given by $*$ extends to a $\mathbf{Q}(v)$ -algebra structure on $\mathfrak{K}(Z_{J,G^0})$ (resp. $\bar{\mathfrak{K}}(Z_{J,G^0})$) denoted again by $*$. Moreover, $\tilde{\mathfrak{K}}(Z_{J,G^0})$ is a two-sided ideal of $\mathfrak{K}(Z_{J,G^0})$ (see 36.11). Note that $\mathfrak{K}(Z_{J,D})^J$ is also a two-sided ideal of $\mathfrak{K}(Z_{J,G^0})$. This follows from the fact that $\mathfrak{e}_{H,J} : \mathfrak{K}(Z_{J,G^0}) \rightarrow \mathfrak{K}(Z_{H,G^0})$ is an algebra homomorphism for any $H \subset J$ (see 36.4(c)). Since the two summands in the right hand side of 36.13(d) are two-sided ideals, they annihilate each other under the product $*$. We see also that the isomorphism 36.13(e) respects the algebra structures.

From 36.9(f) we see that the two summands in the right hand side of 36.13(d) are stable under $\partial : \mathfrak{K}(Z_{J,G^0}) \rightarrow \mathfrak{K}(Z_{J,G^0})$ and from 36.4(h) we have

$$(a) \quad (x * y : \partial(z)) = (\partial(x) : y * z)$$

for $x, y, z \in \mathfrak{K}(Z_{J,G^0})^J$.

Let $\mathcal{W} := \{w \in {}^J\mathbf{W}; \text{Ad}(w)J = J\}$. We show:

$$(b) \quad \mathfrak{K}(Z_{J,G^0})^J = \oplus_{w \in \mathcal{W}} {}^w\mathfrak{K}(Z_{J,G^0})^J$$

where ${}^w\mathfrak{K}(Z_{J,G^0})^J = {}^w\mathfrak{K}(Z_{J,G^0}) \cap \mathfrak{K}(Z_{J,G^0})^J$. Let $x \in \mathfrak{K}(Z_{J,G^0})^J$. By 36.8(b) we can write uniquely $x = \sum_{w \in {}^J\mathbf{W}} x_w$ where $x_w \in {}^w\mathfrak{K}(Z_{J,G^0})$. It is enough to show that $x_w \in \mathfrak{K}(Z_{J,G^0})^J$ (that is, $(y : x_w) = 0$ for any $y \in \tilde{\mathfrak{K}}(Z_{J,G^0})$) for all w

and $x_w = 0$ (that is, $(y' : x_w) = 0$ for any $y' \in \underline{\mathfrak{K}}(Z_{J,G^0})$) if $\text{Ad}(w)J \neq J$. For x_w, y as above we have $y = \sum_{w' \in {}^J\mathbf{W}} y_{w'}$ with $y_{w'} \in {}^{w'}\underline{\mathfrak{K}}(Z_{J,G^0} \cap \tilde{\mathfrak{K}}(Z_{J,G^0}))$, see 36.10(a). Using 36.9(j) twice, we have $(y : x_w) = (y_w : x_w) = (y_w : x) = 0$, as required. Now assume that $w \in {}^J\mathbf{W}$, $\text{Ad}(w)J \neq J$ and $y' \in \underline{\mathfrak{K}}(Z_{J,G^0})$. By 36.8(b) we have $y' = \sum_{w' \in {}^J\mathbf{W}} y'_{w'}$ with $y'_{w'} \in {}^{w'}\underline{\mathfrak{K}}(Z_{J,G^0})$; moreover, by 36.10(a) we have $y'_{w'} \in \tilde{\mathfrak{K}}(Z_{J,G^0})$. Using 36.9(j) twice, we have $(y' : x_w) = (y'_{w'} : x_w) = (y'_{w'} : x) = 0$, as required. This proves (b).

From definitions we see that

(c) *the decomposition (b) corresponds under 36.13(e) to the decomposition $\bar{\mathfrak{K}}(Z_{J,G^0}) = \oplus_{w \in \mathcal{W}} {}^w\bar{\mathfrak{K}}(Z_{J,G^0})$ (see 36.10(d)).*

Under the isomorphism 36.13(e), the involution $\tilde{\mathfrak{D}} : \bar{\mathfrak{K}}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D})$ corresponds to an involution $\tilde{\mathfrak{D}}' : \underline{\mathfrak{K}}(Z_{J,D})^J \rightarrow \underline{\mathfrak{K}}(Z_{J,D})^J$; this is related to the involution $\tilde{\mathfrak{D}} : \underline{\mathfrak{K}}(Z_{J,D}) \rightarrow \underline{\mathfrak{K}}(Z_{J,D})$ by $\tilde{\mathfrak{D}}'x = \tilde{\mathfrak{D}}x \pmod{\tilde{\mathfrak{K}}(Z_{J,D})}$ for $x \in \underline{\mathfrak{K}}(Z_{J,D})^J$. Hence for $x, x' \in \underline{\mathfrak{K}}(Z_{J,D})^J$ we have

$$(\tilde{\mathfrak{D}}'x : x') = (\tilde{\mathfrak{D}}x : x').$$

Using this and 36.9(h) we see that:

(d) *if $x \in \underline{\mathfrak{K}}(Z_{J,D})^J$ satisfies $(x : \tilde{\mathfrak{D}}x) = 0$ then $x = 0$.*

Writing $\heartsuit$ instead of $\{\bar{\mathbf{Q}}_l\} \subset \underline{\mathfrak{g}}(\mathbf{T})$ we see that $\underline{\mathfrak{K}}^\heartsuit(Z_{J,G^0})$ is a subalgebra of $\underline{\mathfrak{K}}(Z_{J,G^0})$. Hence

$$\underline{\mathfrak{K}}^J := \underline{\mathfrak{K}}^\heartsuit(Z_{J,D}) \cap \underline{\mathfrak{K}}(Z_{J,D})^J$$

is a subalgebra of $\underline{\mathfrak{K}}(Z_{J,D})^J$. From (b) and 36.8(d) we see that

$$(e) \quad \underline{\mathfrak{K}}^J = \oplus_{w \in \mathcal{W}} {}^w\underline{\mathfrak{K}}^J$$

where ${}^w\underline{\mathfrak{K}}^J = {}^w\underline{\mathfrak{K}}^\heartsuit(Z_{J,G^0}) \cap \underline{\mathfrak{K}}^J$. Now $\partial : \underline{\mathfrak{K}}(Z_{J,G^0})^J \rightarrow \underline{\mathfrak{K}}(Z_{J,G^0})^J$ leaves $\underline{\mathfrak{K}}^J$ stable. Moreover, for $w \in \mathcal{W}$ we have

$$(f) \quad \partial({}^w\underline{\mathfrak{K}}^J) = {}^{w^{-1}}\underline{\mathfrak{K}}^J.$$

(see 36.9(l)).

Note that $\mathcal{W}$ is the same as the set of all $w \in \mathbf{W}$ such that the corresponding permutation of the set of roots leaves stable the set of simple roots corresponding to elements of J . Hence $\mathcal{W}$ is a subgroup of $\mathbf{W}$.

36.15. By 36.4(f) we have $(x : x') = (x' : x) = \tau(x * \partial(x'))$ for $x, x' \in \underline{\mathfrak{K}}^J$. By 36.13(b), the bilinear form $(:)$ is non-degenerate on (the finite dimensional vector space) $\underline{\mathfrak{K}}^J$. Hence there is a unique vector $x_0 \in \underline{\mathfrak{K}}^J$ such that $(x_0 : x) = \tau(x)$ for all $x \in \underline{\mathfrak{K}}^J$. Hence for $x, x' \in \underline{\mathfrak{K}}^J$ we have $(x_0 : x * \partial(x')) = \tau(x * \partial(x')) = (x : x')$. Using 36.14(a) we rewrite this as $(\partial(x_0) * x : x') = (x : x')$. (We use also that $\partial^2 = 1$ on $\underline{\mathfrak{K}}^J$.) Using the non-degeneracy of $(:)$ on $\underline{\mathfrak{K}}^J$ we deduce $\partial(x_0) * x = x$ for all $x \in \underline{\mathfrak{K}}^J$. For $x, x' \in \underline{\mathfrak{K}}^J$ we have also $(\partial(x) * x' : x_0) = \tau(\partial(x) * x') = (\partial(x) : \partial(x')) = (x : x')$ (we use 36.4(g)). Using 36.14(a) we rewrite this as

$(x : x' * \partial(x_0)) = (x : x')$. Using the non-degeneracy of $(:)$ on $\underline{\mathfrak{K}}^J$ we deduce $x' * \partial(x_0) = x'$ for all $x' \in \underline{\mathfrak{K}}^J$. We see that the algebra $\underline{\mathfrak{K}}^J$ has a unit element, namely $\mathbf{1} = \partial(x_0)$. By 36.14(e) we have $x_0 = \sum_{w \in \mathcal{W}} x_0^w$ where $x_0^w \in {}^w \underline{\mathfrak{K}}^J$. For any $x \in \underline{\mathfrak{K}}^J$ we have similarly $x = \sum_{w \in \mathcal{W}} x^w$ where $x^w \in {}^w \underline{\mathfrak{K}}^J$. From 36.9(k) we see that $\tau(x) = \tau(x^1)$. Hence $(x_0 : x^1) = \tau(x^1) = \tau(x) = (x_0 : x)$. Using 36.9(j) we see that $(x_0 : x^1) = (x_0^1 : x^1) = (x_0^1 : x)$ hence $(x_0^1 : x) = (x_0 : x)$. Using the non-degeneracy of $(:)$ on $\underline{\mathfrak{K}}^J$ we deduce $x_0 = x_0^1$ that is $x_0 \in {}^1 \underline{\mathfrak{K}}^J$. Using 36.14(f) we deduce that $\mathbf{1} \in {}^1 \underline{\mathfrak{K}}^J$.

36.16. We preserve the setup of 36.14, 36.15. In this subsection we assume that $G = G^0$ is a symplectic group $Sp_{2n}(\mathbf{k})$ ($n \geq 1$) and we describe in this case the structure of the algebra $\underline{\mathfrak{K}}^J$. (The proofs, which depend on results in this and future Sections, will be given elsewhere.) We have $\mathbf{I} = \{s_1, s_2, \dots, s_n\}$ where $s_i s_{i+1}$ has order 3 if $i = 1, 2, \dots, n-2$ and order 4 if $i = n-1$; we have $s_i s_j = s_j s_i$ if $|i - j| \geq 2$.

(i) We have $\underline{\mathfrak{K}}^J = 0$ unless

(*) $J = \{s_{k+1}, s_{k+2}, \dots, s_n\}$ with $0 \leq k \leq n$ such that $n - k = a^2 + a$ for some $a \in \mathbf{N}$.

Now assume that (*) holds. Then $\mathcal{W}$ is a Weyl group of type B_k with standard generators $\sigma_1, \sigma_2, \dots, \sigma_k$ where $\sigma_i = s_i$ for $1 \leq i < k$ and σ_k is the unique element in the subgroup of $\mathbf{W}$ generated by $s_k, s_{k+1}, \dots, s_n$ such that $\sigma_k \in \mathcal{W} - \{1\}$ (if $k \geq 1$). Let $\tilde{l} : \mathcal{W} \rightarrow \mathbf{N}$ be the length function of the Weyl group $\mathcal{W}$.

(ii) For any $w \in \mathcal{W}$ we have $\dim({}^w \underline{\mathfrak{K}}^J) = 1$.

(iii) For any $i \in [1, k]$ there is a unique element $x \in {}^{\sigma_i} \underline{\mathfrak{K}}^J - \{0\}$ such that $(x + \mathbf{1}) * (x - c\mathbf{1}) = 0$ for some $c \in v\mathbf{Z}[v]$; in fact we have $c = v^2$ if $1 \leq i < k$ and $c = v^{4a+2}$ if $i = k$. We set $x = t_i$.

(iv) For any $w \in \mathcal{W}$ there is a unique element $t_w \in {}^w \underline{\mathfrak{K}}^J - \{0\}$ such that the following hold: $t_{\sigma_i} = t_i$ for $i \in [1, k]$; $t_w * t_{w'} = t_{w'w}$ if $w, w' \in \mathcal{W}$, $\tilde{l}(w'w) = \tilde{l}(w) + \tilde{l}(w')$.

We see that $\underline{\mathfrak{K}}^J$ is an Iwahori-Hecke algebra with not necessarily equal parameters. Similar results hold for other classical groups.

37. A MACKEY TYPE FORMULA

37.1. We fix a connected component D of G . With notation in 26.1, if $J, J' \subset \mathbf{I}$, $P \in \mathcal{P}_J, Q \in \mathcal{P}_{J'}, u = \text{pos}(P, Q) \in \mathbf{W}$, then $u \in {}^J \mathbf{W}^{J'}$. Setting $P^Q = (P \cap Q)U_P$, we have $P^Q \in \mathcal{P}_{J \cap \text{Ad}(u)J'}$.

For $K, K' \subset \mathbf{I}$ and $u \in {}^K \mathbf{W}^{K'}$ let

$$\Upsilon_u = \{(X, Y, g(U_X \cap U_Y)); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\ g(U_X \cap U_Y) \in D/(U_X \cap U_Y), \text{pos}(X, Y) = u\}.$$

We have a diagram

$$Z_{K \cap \text{Ad}(u)K', D} \xleftarrow{j} \Upsilon_u \xrightarrow{h} Z_{K' \cap \text{Ad}(u^{-1})K, D}$$

where $j(X, Y, g(U_X \cap U_Y)) = (X, gU_X)$, $h(X, Y, g(U_X \cap U_Y)) = (Y, gU_Y)$. Set

$$\Phi_u = h!j^* : \mathcal{D}(Z_{K \cap \text{Ad}(u)K', D}) \rightarrow \mathcal{D}(Z_{K' \cap \text{Ad}(u^{-1})K, D}).$$

Proposition 37.2. *Let $K, K', J \subset \mathbf{I}$ be such that $K \subset J, K' \subset J$. Let $A' \in \mathcal{D}(Z_{K, D})$. We set $\mathfrak{B} = \mathfrak{e}_{K', J} \mathfrak{f}_{K, J} A' \in \mathcal{D}(Z_{K', D})$. For any $u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J$ we set $\mathfrak{B}_u = \mathfrak{f}_{K' \cap \text{Ad}(u^{-1})K, K'} \Phi_u \mathfrak{e}_{K \cap \text{Ad}(u)K', K} A' \in \mathcal{D}(Z_{K', D})$ and $m_u = \dim(U_P \cap U_R)/U_Q$ where $P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}$, $\text{pos}(P, R) = u$ and $Q = Q_{J, P} = Q_{J, R} \in \mathcal{P}_J$ (notation of 36.4). We have*

$$\mathfrak{B} \simeq \{\mathfrak{B}_u[[-m_u]]; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J\},$$

with $\simeq$ as in 32.15.

We have a commutative diagram with a cartesian square

$$\begin{array}{ccccc} \mathfrak{E} & \xrightarrow{\mathfrak{b}} & Z_{K', J, D} & \xrightarrow{\mathfrak{c}'} & Z_{K', D} \\ \mathfrak{a} \downarrow & & \mathfrak{d}' \downarrow & & \\ Z_{K, D} & \xleftarrow{\mathfrak{c}} & Z_{K, J, D} & \xrightarrow{\mathfrak{d}} & Z_{J, D} \end{array}$$

Here

$$\begin{aligned} \mathfrak{E} &= \{(P, R, gU_Q); P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}, gU_Q \in D/U_Q, Q = Q_{J, P} = Q_{J, R}\}, \\ \mathfrak{c}(P, gU_Q) &= (P, gU_P), \mathfrak{d}(P, gU_Q) = (Q, gU_Q) \text{ with } Q = Q_{J, P}, \\ \mathfrak{c}'(R, gU_Q) &= (R, gU_R), \mathfrak{d}'(R, gU_Q) = (Q, gU_Q) \text{ with } Q = Q_{J, R}, \\ \mathfrak{a}(P, R, gU_Q) &= (P, gU_Q), \mathfrak{b}(P, R, gU_Q) = (R, gU_Q). \end{aligned}$$

We have

$$\mathfrak{B} = \mathfrak{c}'_! \mathfrak{d}'^* \mathfrak{d}_! \mathfrak{c}^* A' = \mathfrak{c}'_! \mathfrak{b}_! \mathfrak{a}^* \mathfrak{c}^* A' = (\mathfrak{c}' \mathfrak{b})_! (\mathfrak{c} \mathfrak{a})^* A' = \mathfrak{q}_! \mathfrak{p}^* A'$$

where $\mathfrak{q} = \mathfrak{c}' \mathfrak{b} : \mathfrak{E} \rightarrow Z_{K', D}$, $\mathfrak{p} = \mathfrak{c} \mathfrak{a} : \mathfrak{E} \rightarrow Z_{K, D}$ are given by $\mathfrak{q}(P, R, gU_Q) = (R, gU_R)$, $\mathfrak{p}(P, R, gU_Q) = (P, gU_P)$. We have a partition

$$\mathfrak{E} = \sqcup_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J} \mathfrak{E}_u$$

where $\mathfrak{E}_u = \{(P, R, gU_Q) \in \mathfrak{E}; \text{pos}(P, R) = u\}$ is locally closed in $\mathfrak{E}$. Let $\mathfrak{p}_u = \mathfrak{p}|_{\mathfrak{E}_u} : \mathfrak{E}_u \rightarrow Z_{K, D}$, $\mathfrak{q}_u = \mathfrak{q}|_{\mathfrak{E}_u} : \mathfrak{E}_u \rightarrow Z_{K'}$. By 32.15, we have

$$\mathfrak{q}_! \mathfrak{p}^* A' \simeq \{\mathfrak{q}_u! \mathfrak{p}_u^* A'; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J\}.$$

It remains to show that, for u as above, we have

$$\mathfrak{q}_u! \mathfrak{p}_u^* A' = \mathfrak{B}_u[[-m_u]].$$

We have a commutative diagram

$$\begin{array}{ccccc}
Z_{K,D} & \xleftarrow{\tilde{\mathfrak{d}}} & Z_{K \cap \text{Ad}(u)K', K, D} & \xrightarrow{\tilde{\mathfrak{c}}} & Z_{K \cap \text{Ad}(u)K', D} \\
& & \uparrow \tilde{\mathfrak{a}} & & \uparrow \mathfrak{j} \\
& & \mathfrak{S}'_u & \xrightarrow{\tilde{\mathfrak{b}}} & \Upsilon_u & \xrightarrow{\mathfrak{h}} & Z_{K' \cap \text{Ad}(u^{-1})K, D} \\
& & \uparrow \mathfrak{r} & & & & \uparrow \tilde{\mathfrak{c}}' \\
& & \mathfrak{S}_u & \xrightarrow{\mathfrak{y}} & & \longrightarrow & Z_{K' \cap \text{Ad}(u^{-1})K, K', D} \\
& & & & & & \downarrow \tilde{\mathfrak{d}}' \\
& & & & & & Z_{K', D}
\end{array}$$

Here

$$\begin{aligned}
& \tilde{\mathfrak{c}}(X, gU_P) = (X, gU_X), \tilde{\mathfrak{d}}(X, gU_P) = (P, gU_P), \\
& \tilde{\mathfrak{c}}'(Y, gU_R) = (Y, gU_Y), \tilde{\mathfrak{d}}'(Y, gU_R) = (R, gU_R), \\
& \mathfrak{S}'_u, \tilde{\mathfrak{a}}, \tilde{\mathfrak{b}} \text{ are defined so that the square } (\tilde{\mathfrak{a}}, \tilde{\mathfrak{b}}, \tilde{\mathfrak{c}}, \mathfrak{j}) \text{ is cartesian;} \\
& \mathfrak{S}_u, \mathfrak{r}, \mathfrak{y} \text{ are defined so that the square } (\mathfrak{r}, \mathfrak{y}, \tilde{\mathfrak{b}}, \tilde{\mathfrak{c}}') \text{ is cartesian.}
\end{aligned}$$

Then

$$\begin{aligned}
\mathfrak{S}_u = \{ & (X, Y, g(U_X \cap U_Y), g'U_P, g''U_R); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\
& g(U_X \cap U_Y) \in D/(U_X \cap U_Y), g'U_P \in D/U_P, g''U_R \in D/U_R, \text{pos}(X, Y) = u \\
& , P = Q_{K, X}, R = Q_{K', Y}, g'U_X = gU_X, g''U_Y = gU_Y \}.
\end{aligned}$$

Set $\mathfrak{r} = \tilde{\mathfrak{d}}'\mathfrak{y} : \mathfrak{S}_u \rightarrow Z_{K', D}$, $\mathfrak{s} = \tilde{\mathfrak{d}}\tilde{\mathfrak{a}}\mathfrak{r} : \mathfrak{S}_u \rightarrow Z_{K, D}$. Then

$$\begin{aligned}
& \mathfrak{r}(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) = (P, g'U_P), \\
& \mathfrak{s}(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) = (R, g''U_R).
\end{aligned}$$

We have

$$\begin{aligned}
\mathfrak{B}_u &= \tilde{\mathfrak{d}}'\tilde{\mathfrak{c}}'^*\mathfrak{h}_!\mathfrak{j}^*\tilde{\mathfrak{c}}_!\tilde{\mathfrak{d}}^*A' = \tilde{\mathfrak{d}}'\tilde{\mathfrak{c}}'^*\mathfrak{h}_!\tilde{\mathfrak{b}}_!\tilde{\mathfrak{a}}^*\tilde{\mathfrak{d}}^*A' \\
&= \tilde{\mathfrak{d}}'\tilde{\mathfrak{c}}'^*(\mathfrak{h}\tilde{\mathfrak{b}})_!(\tilde{\mathfrak{d}}\tilde{\mathfrak{a}})^*A' = \tilde{\mathfrak{d}}'\mathfrak{y}_!\mathfrak{r}^*(\tilde{\mathfrak{d}}\tilde{\mathfrak{a}})^*A' = (\tilde{\mathfrak{d}}'\mathfrak{y})_!(\tilde{\mathfrak{d}}\tilde{\mathfrak{a}}\mathfrak{r})^*A' = \mathfrak{r}_!\mathfrak{s}^*A'.
\end{aligned}$$

We show that $\mathfrak{q}_u!\mathfrak{p}_u^*A' = \mathfrak{r}_!\mathfrak{s}^*A'[[-m_u]]$. We have a commutative diagram

$$\begin{array}{ccccc}
Z_{K,D} & \xleftarrow{\mathfrak{p}_u} & \mathfrak{E}_u & \xrightarrow{\mathfrak{q}_u} & Z_{K', D} \\
1 \downarrow & & \downarrow \mathfrak{t} & & 1 \downarrow \\
Z_{K,D} & \xleftarrow{\mathfrak{s}} & \mathfrak{S}_u & \xrightarrow{\mathfrak{r}} & Z_{K', D}
\end{array}$$

where $\mathbf{t}(P, R, gU_Q) = (P^R, R^P, g(U_{P^R} \cap U_{R^P}), gU_P, gU_R)$ is well defined since $U_Q \subset U_P, U_Q \subset U_R, U_Q \subset U_{P^R}, U_Q \subset U_{R^P}$. We continue the proof assuming that

(a) $\mathbf{t}$ is an affine space bundle with fibres of dimension m_u .

For any $\tilde{A} \in \mathcal{D}(\mathfrak{S}_u)$ we have $\mathbf{t}_! \mathbf{t}^*(\tilde{A}) = \tilde{A}[[-m_u]]$. Hence

$$\mathbf{r}_! \mathbf{s}^* A'[[-m_u]] = \mathbf{r}_! \mathbf{t}_! \mathbf{t}^* \mathbf{s}^* A' = (\mathbf{r}\mathbf{t})_! (\mathbf{s}\mathbf{t})^* A' = \mathbf{q}_u! \mathbf{p}_u^* A',$$

as required.

We prove (a). We only show that each fibre of $\mathbf{t}$ is an affine space of dimension m_u . First we show that $\mathbf{t} : \mathfrak{E}_u \rightarrow \mathfrak{S}_u$ is surjective. Assume that we are given $(X, Y, g_0(U_X \cap U_Y), g'_0 U_P, g''_0 U_R) \in \mathfrak{S}_u$. Then $g_0, g'_0, g''_0 \in D$, $v' = g_0^{-1} g'_0 \in U_X$, $v'' = g_0^{-1} g''_0 \in U_Y$. We must show that there exists $g \in D$ such that $g_0 \in g(U_X \cap U_Y)$, $g'_0 \in gU_P$, $g''_0 \in gU_R$. Setting $y = g^{-1} g_0$, we must show that there exists $y \in U_X \cap U_Y$ such that $yv' \in U_P, yv'' \in U_R$. We have $v' = v'_1 v'_2$ where $v'_1 \in U_R \cap P$, $v'_2 \in U_P$ and $v'' = v''_1 v''_2$ where $v''_1 \in U_P \cap R$, $v''_2 \in U_R$. Then $v'_1 \in U_X \cap U_Y, v''_1 \in U_X \cap U_Y$. Setting $y = (v'_1 v''_1)^{-1} \in U_X \cap U_Y$ we have $yv' = v''_1^{-1} v'_2 \in U_P$ and $yv'' = v''_1^{-1} v'_1^{-1} v''_2 \in v''_1^{-1} U_R v''_1 U_R = U_R$, as desired.

It remains to show that, if $(P, R, gU_Q) \in \mathfrak{E}_u$, then

(b) $F = \{(P', R', g'U_{Q'}) \in E_u; \mathbf{t}(P', R', g'U_{Q'}) = \mathbf{t}(P, R, gU_Q)\}$ is an affine space of dimension m_u .

For $(P', R', g'U_{Q'}) \in F$, both P, P' contain $P^R = (P')^{R'}$ and have the same type hence $P = P'$. Similarly $R = R', Q = Q'$. Hence

$$\begin{aligned} F &\cong \{g'U_Q; gU_P = g'U_P, gU_R = g'U_R, g(U_{P^R} \cap U_{R^P}) = g'(U_{P^R} \cap U_{R^P})\} \\ &= \{g'U_Q; g^{-1}g' \in U_P \cap U_R \cap U_{P^R} \cap U_{R^P}\} \\ &= \{g'U_Q; g^{-1}g' \in U_P \cap U_R\} \cong (U_P \cap U_R)/U_Q, \end{aligned}$$

and (b) follows. This completes the proof.

37.3. In the remainder of this section we assume that we are in the setup of 36.8. From 37.2 we deduce that for $K, K', J \subset \mathbf{I}$ such that $K \subset J, K' \subset J$ we have

$$\mathfrak{e}_{K', J} \mathfrak{f}_{K, J} = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J} v^{m_u} \mathfrak{f}_{K' \cap \text{Ad}(u^{-1})K, K'} \Phi_u \mathfrak{e}_{K \cap \text{Ad}(u)K', K}$$

as $\mathcal{A}$ -linear maps $\mathfrak{K}(Z_{K, D}) \rightarrow \mathfrak{K}(Z_{K', D})$, where the $\mathcal{A}$ -linear map

$\mathfrak{K}(Z_{K \cap \text{Ad}(u)K', D}) \rightarrow \mathfrak{K}(Z_{K' \cap \text{Ad}(u^{-1})K, D})$
induced by Φ_u in 37.1 is denoted again by Φ_u .

37.4. Let $K, K' \subset J$ and let $x \in \mathfrak{K}(Z_{K, D})^K, x' \in \mathfrak{K}(Z_{K', D})^{K'}$. We show:

$$(a) \quad (\mathfrak{f}_{K, J} x : \mathfrak{f}_{K', J} x') = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J; K = \text{Ad}(u)K'} v^{m(u)} (\Phi_u x : x').$$

Here $m(u)$, Φ_u are as in 37.3; in our case, $\Phi_u : \mathfrak{K}(Z_{K,D}) \rightarrow \mathfrak{K}(Z_{K',D})$. Using 37.3 we have

$$\begin{aligned}
(\mathfrak{f}_{K,J}x : \mathfrak{f}_{K',J}x') &= (\mathfrak{e}_{K',J}\mathfrak{f}_{K,J}x : x') \\
&= \left(\sum_{u \in {}^K\mathbf{W}^{K'} \cap \mathbf{W}_J} v^{m_u} \mathfrak{f}_{K' \cap \text{Ad}(u^{-1})K, K'} \Phi_u \mathfrak{e}_{K \cap \text{Ad}(u)K', K} x : x' \right) \\
&= \sum_{u \in {}^K\mathbf{W}^{K'} \cap \mathbf{W}_J} v^{m_u} (\Phi_u \mathfrak{e}_{K \cap \text{Ad}(u)K', K} x : \mathfrak{e}_{K' \cap \text{Ad}(u^{-1})K, K'} x') \\
&= \sum_{u \in {}^K\mathbf{W}^{K'} \cap \mathbf{W}_J; K \cap \text{Ad}(u)K' = K, K' \cap \text{Ad}(u^{-1})K = K'} v^{m_u} (\Phi_u x : x').
\end{aligned}$$

The condition that $K \cap \text{Ad}(u)K' = K, K' \cap \text{Ad}(u^{-1})K = K'$ is equivalent to $K \subset \text{Ad}(u)K', K' \subset \text{Ad}(u^{-1})K$ that is, to $K = \text{Ad}(u)K'$. This proves (a).

Consider the equivalence relation on the set of subsets of J given by $K_1 \sim K_2$ if $\text{Ad}(u)K_1 = K_2$ for some $u \in \mathbf{W}_J$. For any equivalence class $\mathfrak{o}$ under $\sim$ we set

$$\underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o} = \sum_{H \subset J; H \in \mathfrak{o}} f_{H,J}(\underline{\mathfrak{K}}(Z_{H,D})^H) \subset \underline{\mathfrak{K}}(Z_{J,D}).$$

Note that $\underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o}$ is homogeneous.

Proposition 37.5. *We have*

$$(a) \quad \underline{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\mathfrak{o}} \underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o}$$

where $\mathfrak{o}$ runs over the equivalence classes for $\sim$.

If $J = \emptyset$ we have $\underline{\mathfrak{K}}(Z_{J,D}) = \underline{\mathfrak{K}}(Z_{J,D})^J$ and the result is obvious. We may assume that $J \neq \emptyset$ and that the result is true when J is replaced by a strictly smaller subset. Using 36.13(d) and the induction hypothesis we have

$$\begin{aligned}
\underline{\mathfrak{K}}(Z_{J,D}) &= \underline{\mathfrak{K}}(Z_{J,D})^J + \sum_{L \subsetneq J} \mathfrak{f}_{L,J} \sum_{J'; J' \subset L} \mathfrak{f}_{J',L} \underline{\mathfrak{K}}(Z_{J',D})^{J'} \\
&\subset \sum_{J'; J' \subset J} \mathfrak{f}_{J',J} \underline{\mathfrak{K}}(Z_{J',D})^{J'}.
\end{aligned}$$

Thus, $\underline{\mathfrak{K}}(Z_{J,D}) = \sum_{\mathfrak{o}} \underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o}$. Next we show that $(\underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o} : \underline{\mathfrak{K}}(Z_{J,D})^{\mathfrak{o}'}) = 0$ if $\mathfrak{o} \neq \mathfrak{o}'$. It is enough to show that $(\mathfrak{f}_{H,J}(\underline{\mathfrak{K}}(Z_{H,D})^H) : \mathfrak{f}_{H',J}(\underline{\mathfrak{K}}(Z_{H',D})^{H'})) = 0$ if $H, H' \subset J, H \not\sim H'$. This follows from 37.4(a). It remains to use the following (easily verified) statement: if V is a finite dimensional vector space with a non-singular symmetric bilinear form $(:)$ and $V_1, V_2, \dots, V_k$ are subspaces such that $(V_i : V_j) = 0$ for $i \neq j$ and $\sum_i V_i = V$ then $V = \bigoplus_i V_i$.

38. DUALITY

38.1. We fix a connected component D of G that generates G . In this section we study an involution of the set of isomorphism classes of character sheaves on D called *duality*.

We write ϵ instead of $\epsilon_D : \mathbf{W} \rightarrow \mathbf{W}$ (see 26.2). For $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ we set, as in 30.3:

$$V_{J,D} = \{(P, gU_P); P \in \mathcal{P}_J, gU_P \in N_D P/U_P\}.$$

We also set $\mathbf{W}_J^\epsilon = \{w \in \mathbf{W}_J; \epsilon(w) = w\}$ where $\mathbf{W}_J$ is as in 26.1. Let J_ϵ be the set of orbits of the restriction of ϵ to J .

For any $P_0 \in \mathcal{P}_J$ we have a functor $A_0 \mapsto A_0^\flat$ (see 30.3) from the category of P_0/U_{P_0} -equivariant perverse sheaves on the connected component $N_D P_0/U_{P_0}$ of $N_G P_0/U_{P_0}$ to the category of perverse sheaves on $V_{J,D}$.

Let $CS(V_{J,D})$ be the full subcategory of the category of perverse sheaves on $V_{J,D}$ whose objects are isomorphic to objects of the form $A_0^\flat$ where A_0 is a direct sum of character sheaves on $N_D P_0/U_{P_0}$. In particular, $CS(V_{\mathbf{I},D})$ (denoted also by $CS(D)$) is the category of perverse sheaves on D that are direct sums of character sheaves. Note that $A_0 \mapsto A_0^\flat$ is an equivalence of categories $CS(N_D P_0/U_{P_0}) \xrightarrow{\sim} CS(V_{J,D})$.

For $J \subset J' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J'$ we have functors $f_{J,J'} : \mathcal{D}(V_{J,D}) \rightarrow \mathcal{D}(V_{J',D})$ and $e_{J,J'} : \mathcal{D}(V_{J',D}) \rightarrow \mathcal{D}(V_{J,D})$, see 30.4. From definitions, for $J \subset J' \subset J'' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J', \epsilon(J'') = J''$, we have

$$(a) \quad f_{J,J''} = f_{J',J''} f_{J,J'}, \quad e_{J,J''} = e_{J,J'} e_{J',J''}.$$

Clearly, $f_{J,J} = 1, e_{J,J} = 1$.

38.2. We show that:

(a) for $J \subset J'$ as above, $e_{J,J'}$ restricts to a functor $CS(V_{J',D}) \rightarrow CS(V_{J,D})$ denoted again by $e_{J,J'}$.

Let $P \in \mathcal{P}_J, P' \in \mathcal{P}_{J'}$ be such that $P \subset P'$. Let $D_0 = N_D P/U_P, D'_0 = N_D P'/U_{P'}$. Let $C_0 \in CS(D'_0)$ and let $C = C_0^\flat$ be the corresponding object of $CS(V_{J',D})$. From 31.14 we see that $\text{res}_{D'_0}^{D_0}(C_0) \in CS(D_0)$ and from 30.4(b) we see that $e_{J,J'} C$ is the perverse sheaf $(\text{res}_{D'_0}^{D_0}(C_0))^\flat$ on $V_{J,D}$ hence $e_{J,J'} C \in CS(V_{J,D})$. This proves (a).

38.3. We show that:

(a) for $J \subset J'$ as above, $f_{J,J'}$ restricts to a functor $CS(V_{J,D}) \rightarrow CS(V_{J',D})$ denoted again by $f_{J,J'}$.

Let P, P', D_0, D'_0 be as in 38.2. Let $A_0 \in CS(D_0)$ and let $A = A_0^\flat$ be the corresponding object of $CS(V_{J,D})$. By 30.4(a), $f_{J,J'} A = A'_0{}^\flat$ where $A'_0 = \text{ind}_{D_0}^{D'_0} A_0$ is a direct sum of simple admissible perverse sheaves on D'_0 . It remains to show that $A'_0 \in CS(D'_0)$. To do this we may assume that $J' = \mathbf{I}$ hence $P' = G^0, D'_0 = D$. Let $\alpha = \dim U_P$. Let L be a Levi of P . We can identify naturally $N_G P/U_P$ with

$H = N_G P \cap N_G L$, a reductive group with $H^0 = L$. Then D_0 becomes $H \cap D$. We identify the canonical torus of L with the canonical torus $\mathbf{T}$ of G^0 , and the Weyl group of L with the subgroup $\mathbf{W}_J$ of $\mathbf{W}$ as in 29.1. Let $\mathcal{L} \in \mathfrak{s}(\mathbf{T})$. Let $\mathbf{s} = (s_1, s_2, \dots, s_r)$ be a sequence in $J \cup \{1\}$ such that $s_1 s_2 \dots s_r \underline{D} \in \mathbf{W}_{\mathcal{L}}^{\bullet}$ (see 28.3). Let $\bar{Z}_{\emptyset, \mathbf{I}, D}^{\mathbf{s}}, \bar{K}_D^{\mathbf{s}, \mathcal{L}} \in \mathcal{D}(D)$ be as in 28.12, $\bar{\mathcal{L}}$ as in 28.9, and let $\bar{Z}_{\emptyset, J, D_0}^{\mathbf{s}}, \bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}} \in \mathcal{D}(D_0)$, $\bar{\mathcal{L}}_0$, be the analogous objects defined in terms of H, D_0 instead of G, D . Consider the commutative diagram

$$\begin{array}{ccccc}
\bar{Z}_{\emptyset, J, D_0}^{\mathbf{s}} & \xleftarrow{pr_2} & V_1 \times_{D_0} \bar{Z}_{\emptyset, J, D_0}^{\mathbf{s}} & \xrightarrow{f_0} & \bar{Z}_{\emptyset, \mathbf{I}, D}^{\mathbf{s}} \\
f_1 \downarrow & & pr_1 \downarrow & & f_2 \downarrow \\
D_0 & \xleftarrow{a_1} & V_1 & \xrightarrow{a'} & V_2 \xrightarrow{a''} D
\end{array}$$

where

$$\begin{aligned}
V_1 &= \{(g, x) \in D \times G^0; x^{-1}gx \in N_D P\}, \\
V_2 &= \{(g, xP) \in D \times G^0/P; x^{-1}gx \in N_D P\}, \\
a_1(g, x) &= g' \text{ where } g_0 \in D_0 \text{ is such that } x^{-1}gx \in g_0 U_P, \\
a'(g, x) &= (g, xP), a''(g, xP) = g, f_1(\beta_0, \beta_1, \dots, \beta_r, g_0) = g_0, \\
f_2(B_0, B_1, \dots, B_r, g) &= (g, x_0 P) \text{ where } x_0 \in G^0 \text{ is such that } x_0^{-1} B_0 x_0 \subset P, \\
f_0((g, x), (\beta_0, \beta_1, \dots, \beta_r, g_0)) &= (x\beta_0 U_P x^{-1}, x\beta_1 U_P x^{-1}, \dots, x\beta_r U_P x^{-1}, g).
\end{aligned}$$

Both squares in the diagram are cartesian and the maps a', f_0 (resp. a_1, pr_2) are smooth with connected fibres of dimension $\dim G - \alpha$ (resp. $\dim G + \alpha$). From definitions we have $pr_2^* \bar{\mathcal{L}}' = f_0^* \bar{\mathcal{L}}$. Hence

$$a_1^*(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}}) = a_1^* f_1^* \bar{\mathcal{L}}_0 = pr_{1!} pr_2^* \bar{\mathcal{L}}_0 = pr_{1!} f_0^* \bar{\mathcal{L}} = a'^* f_2^* \bar{\mathcal{L}}.$$

We see that

$$a_1^{\star}(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}})[- \dim G - \alpha] = a'^{\star} \tilde{K}[- \dim G + \alpha]$$

where $\tilde{K} = f_2^* \bar{\mathcal{L}}$, that is $a_1^{\star}(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}}) = a'^{\star} \tilde{K}[2\alpha]$. Hence

$$\begin{aligned}
a_1^{\star}(^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}})) &= ^p H^i(a_1^{\star}(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}})) = ^p H^i(a'^{\star} \tilde{K}[2\alpha]) = a'^{\star}(^p H^i(\tilde{K}[2\alpha])) \\
&= a'^{\star}(^p H^{i+2\alpha} \tilde{K}).
\end{aligned}$$

From this and definition (27.1) we see that

$$\text{ind}_{D_0}^D(^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}})) = a_1''(^p H^{i+2\alpha} \tilde{K}).$$

We have

$$(b) \quad \oplus_i \text{ind}_{D_0}^D(^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}}))[-i] = \oplus_i ^p H^{i+2\alpha}(\bar{K}_D^{\mathbf{s}, \mathcal{L}})[-i] \in \mathcal{D}(D).$$

Indeed, the left hand side is

$$\begin{aligned} \oplus_i a_!''({}^p H^{i+2\alpha} \tilde{K})[-i] &= a_!''(\tilde{K}[2\alpha]) = a_!'' f_{2!} \bar{\mathcal{L}}[2\alpha] = \bar{K}_D^{\mathbf{s}, \mathcal{L}}[2\alpha] \\ &= \oplus_i {}^p H^{i+2\alpha}(\bar{K}_D^{\mathbf{s}, \mathcal{L}})[-i] \in \mathcal{D}(D), \end{aligned}$$

where we use that $\tilde{K}$ and $\bar{K}_D^{\mathbf{s}, \mathcal{L}}$ are semisimple complexes (a consequence of the decomposition theorem [BBD]). Since ${}^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}})$ is a direct sum of character sheaves on D_0 , we see, using 30.6(a), that $\text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}}))$ is a perverse sheaf on D . Taking ${}^p H^i$ for both sides of (b) we therefore find

$$(c) \quad \text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}})) = {}^p H^{i+2\alpha}(\bar{K}_D^{\mathbf{s}, \mathcal{L}})$$

for any $i \in \mathbf{Z}$. To prove (a) it is enough to verify the following statement.

(d) *If $A_1 \in \hat{D}_0^{\mathcal{L}}$, then $\text{ind}_{D_0}^D(A_1)$ is a direct sum of character sheaves in $\hat{D}^{\mathcal{L}}$.*

We may assume that A_1 is a direct summand of ${}^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}})$. Then $\text{ind}_{D_0}^D(A_1)$ is a direct summand of $\text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}}))$. From (c) we see that $\text{ind}_{D_0}^D(A_1)$ is a direct summand of ${}^p H^{i+2\alpha}(\bar{K}_D^{\mathbf{s}, \mathcal{L}})$ which is a direct sum of character sheaves in $\hat{D}^{\mathcal{L}}$. This proves (d) hence also (a).

38.4. For $J \subset J' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J'$ we have functors $\tilde{f}_{J, J'} : \mathcal{D}(V_{J, D}) \rightarrow \mathcal{D}(V_{J', D})$, $\tilde{e}_{J, J'} : \mathcal{D}(V_{J', D}) \rightarrow \mathcal{D}(V_{J, D})$, see 30.4.

Let $K, K', J \subset \mathbf{I}$ be such that $K \subset J, K' \subset J, \epsilon(J) = J, \epsilon(K) = K, \epsilon(K') = K'$. For any $u \in {}^K \mathbf{W}^{K'} \in \mathbf{W}_J^\epsilon$ let

$$\begin{aligned} \Xi_u = \{ & (X, Y, g(U_X \cap U_Y)); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\ & g(U_X \cap U_Y) \in (N_D X \cap N_D Y)/(U_X \cap U_Y), \text{pos}(X, Y) = u \}. \end{aligned}$$

We have a diagram

$$V_{K \cap \text{Ad}(u)K', D} \xleftarrow{j} \Xi_u \xrightarrow{h} V_{K' \cap \text{Ad}(u^{-1})K, D}$$

where $j(X, Y, g(U_X \cap U_Y)) = (X, gU_X)$, $h(X, Y, g(U_X \cap U_Y)) = (Y, gU_Y)$. Set $\Psi_u = h_! j^* : \mathcal{D}(V_{K \cap \text{Ad}(u)K', D}) \rightarrow \mathcal{D}(V_{K' \cap \text{Ad}(u^{-1})K, D})$.

Lemma 38.5. *Let $A' \in \mathcal{D}(V_{K, D})$. We set $\tilde{\mathcal{C}} = \tilde{e}_{K', J} \tilde{f}_{K, J} A' \in \mathcal{D}(V_{K', D})$. For any $u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon$ we set $\tilde{\mathcal{C}}_u = \tilde{f}_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi_u \tilde{e}_{K \cap \text{Ad}(u)K', K} A' \in \mathcal{D}(V_{K', D})$ and $m_u = \dim(U_P \cap U_R)/U_Q$ where $P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}, \text{pos}(P, R) = u$ and $Q = Q_{J, P} = Q_{J, R} \in \mathcal{P}_J$ (notation of 36.4). We have*

$$\tilde{\mathcal{C}} \simeq \{ \tilde{\mathcal{C}}_u[[-m_u]]; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon \}$$

where $\simeq$ is as in 32.15.

The proof is very similar to that of Proposition 37.2. We have a commutative diagram with a cartesian square

$$\begin{array}{ccccc}
 E & \xrightarrow{b} & V_{K',J,D} & \xrightarrow{c'} & V_{K',D} \\
 a \downarrow & & & & d' \downarrow \\
 V_{K,D} & \xleftarrow{c} & V_{K,J,D} & \xrightarrow{d} & V_{J,D}
 \end{array}$$

Here

$$E = \{(P, R, gU_Q); P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}, gU_Q \in (N_D P \cap N_D R)/U_Q, Q = Q_{J,P} = Q_{J,R}\},$$

$$c(P, gU_Q) = (P, gU_P), d(P, gU_Q) = (Q, gU_Q) \text{ with } Q = Q_{J,P},$$

$$c'(R, gU_Q) = (R, gU_R), d'(R, gU_Q) = (Q, gU_Q), \text{ with } Q = Q_{J,R},$$

$$a(P, R, gU_Q) = (P, gU_Q), b(P, R, gU_Q) = (R, gU_Q).$$

We have

$$\tilde{\mathfrak{C}} = c'_! d'^* d_! c^* A' = c'_! b_! a^* c^* A' = (c'b)_!(ca)^* A' = q_! p^* A'$$

where $q = c'b : E \rightarrow V_{K',D}$, $p = ca : E \rightarrow V_{K,D}$ are given by $q(P, R, gU_Q) = (R, gU_R)$, $p(P, R, gU_Q) = (P, gU_P)$. We have a partition

$$E = \sqcup_{u \in K} \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon E_u$$

where $E_u = \{(P, R, gU_Q) \in E; \text{pos}(P, R) = u\}$ is locally closed in E . Let $p_u = p|_{E_u} : E_u \rightarrow V_{K,D}$, $q_u = q|_{E_u} : E_u \rightarrow V_{K',D}$. By 32.15, we have

$$q_! p^* A' \simeq \{q_u! p_u^* A'; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}.$$

It remains to show that, for u as above, we have $q_u! p_u^* A' = \tilde{\mathfrak{C}}_u[[-m_u]]$. We have a commutative diagram

$$\begin{array}{ccccccc}
 V_{K,D} & \xleftarrow{\tilde{d}} & V_{K \cap \text{Ad}(u)K', K, D} & \xrightarrow{\tilde{c}} & V_{K \cap \text{Ad}(u)K', D} & & \\
 & & \tilde{a} \uparrow & & j \uparrow & & \\
 & & S'_u & \xrightarrow{\tilde{b}} & \Xi_u & \xrightarrow{h} & V_{K' \cap \text{Ad}(u^{-1})K, D} \\
 & & x \uparrow & & & & \tilde{c}' \uparrow \\
 & & S_u & \xrightarrow{y} & & \longrightarrow & V_{K' \cap \text{Ad}(u^{-1})K, K', D} \\
 & & & & & & \tilde{d}' \downarrow \\
 & & & & & & V_{K', D}
 \end{array}$$

Here

$\tilde{c}(X, gU_P) = (X, gU_X), \tilde{d}(X, gU_P) = (P, gU_P),$
 $\tilde{c}'(Y, gU_R) = (Y, gU_Y), \tilde{d}'(Y, gU_R) = (R, gU_R),$
 $S'_u, \tilde{a}, \tilde{b}$ are defined so that the square $(\tilde{a}, \tilde{b}, \tilde{c}, j)$ is cartesian;
 S_u, x, y are defined so that the square $(x, y, h\tilde{b}, \tilde{c}')$ is cartesian.

Then

$$\begin{aligned}
 S_u &= \{(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\
 &g(U_X \cap U_Y) \in (N_D X \cap N_D Y)/(U_X \cap U_Y), g'U_P \in N_D X/U_P, \\
 &g''U_R \in N_D Y/U_R, \text{pos}(X, Y) = u, P = Q_{K, X}, R = Q_{K', Y}, \\
 &g'U_X = gU_X, g''U_Y = gU_Y\}.
 \end{aligned}$$

Set $r = \tilde{d}'y : S_u \rightarrow V_{K', D}, s = \tilde{d}\tilde{a}x : S_u \rightarrow V_{K, D}$. Then

$$\begin{aligned}
 r(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) &= (P, g'U_P), \\
 s(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) &= (R, g''U_R).
 \end{aligned}$$

We have

$$\begin{aligned}
 \tilde{\mathfrak{C}}_u &= \tilde{d}'_! \tilde{c}'^* h_{!j}^* \tilde{c}_! \tilde{d}^* A' = \tilde{d}'_! \tilde{c}'^* h_{!} \tilde{b}_! \tilde{a}^* \tilde{d}^* A' = \tilde{d}'_! \tilde{c}'^* (h\tilde{b})_! (\tilde{d}\tilde{a})^* A' \\
 &= \tilde{d}'_! y_{!x}^* (\tilde{d}\tilde{a})^* A' = (\tilde{d}'y)_! (\tilde{d}\tilde{a}x)^* A' = r_! s^* A'.
 \end{aligned}$$

We show that $q_{u!} p_u^* A' = r_! s^* A' [[-m_u]]$. We have a commutative diagram

$$\begin{array}{ccccc}
 V_{K, D} & \xleftarrow{p_u} & E_u & \xrightarrow{q_u} & V_{K', D} \\
 1 \downarrow & & t \downarrow & & 1 \downarrow \\
 V_{K, D} & \xleftarrow{s} & S_u & \xrightarrow{r} & V_{K', D}
 \end{array}$$

where $t(P, R, gU_Q) = (P^R, R^P, g(U_{P^R} \cap U_{R^P}), gU_P, gU_R)$ is well defined since $U_Q \subset U_P, U_Q \subset U_R, U_Q \subset U_{P^R}, U_Q \subset U_{R^P}$. We continue the proof assuming that

(a) t is an affine space bundle with fibres of dimension m_u .

For any $\tilde{A} \in \mathcal{D}(S_u)$ we have $t_! t^*(\tilde{A}) = \tilde{A} [[-m_u]]$. Hence

$$r_! s^* A' [[-m_u]] = r_! t_! t^* s^* A' = (rt)_! (st)^* A' = q_{u!} p_u^* A',$$

as required.

We prove (a). Consider the commutative diagram

$$\begin{array}{ccc}
 E_u & \longrightarrow & \mathfrak{E}_u \\
 t \downarrow & & \mathfrak{t} \downarrow \\
 S_u & \longrightarrow & \mathfrak{S}_u
 \end{array}$$

where $\mathfrak{E}_u \xrightarrow{\mathfrak{t}} \mathfrak{S}_u$ is as in the proof of 37.2 and the horizontal maps are the obvious imbeddings. Clearly, this diagram is cartesian. Hence (a) is a consequence of the analogous statement 37.2(a) for $\mathfrak{E}_u \xrightarrow{\mathfrak{t}} \mathfrak{S}_u$. This completes the proof.

38.6. In the setup of 38.4, let $u \in {}^K\mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon$. Let $\delta = \dim(U_{X_u}/(U_{X_u} \cap U_{Y_u})) = \dim(U_{Y_u}/(U_{X_u} \cap U_{Y_u}))$ where $(X_u, Y_u, g(U_{X_u} \cap U_{Y_u})) \in \Xi_u$. Let $\alpha_u = \dim U_{X_u} = \dim U_{Y_u}$, $\alpha = \dim U_P$, $\alpha' = \dim U_R$ where $P \in \mathcal{P}_K$, $R \in \mathcal{P}_{K'}$. We show:

(a) h and j in 38.4 are affine space bundles with fibres of dimension δ .

It is enough to prove the statements on j (the statement on h is entirely similar). We show that j is surjective. Let X, g be such that $(X, gU_X) \in V_{K \cap \text{Ad}(u)K', D}$. We must show that there exist $Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}$, $g' \in N_D X \cap N_D Y$ such that $\text{pos}(X, Y) = u$ and $g'U_X = gU_X$. Setting $g^{-1}g' = v$, it is enough to show that for any $Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}$ with $\text{pos}(X, Y) = u$ there exists $v \in U_X$ such that $gv \in N_G Y$. Now X, Y contain a common Levi M . Since $g \in N_G X$ we can find $v \in U_X$ such that $g' = gv \in N_G X \cap N_G M$. There is a unique parabolic Y' of the same type as Y such that Y' has Levi M and $\text{pos}(X, Y') = u$. Then $\text{pos}(g'Xg'^{-1}, g'Yg'^{-1}) = u$, $g'Yg'^{-1}$ has Levi $g'Mg'^{-1} = M$. By uniqueness, we have $Y' = Y$. Thus $g' \in N_G Y$.

We show that the fibres of j are affine spaces of dimension δ . Let $(X, Y, g(U_X \cap U_Y)) \in \Xi_u$. We must show that $F = \{(X, Y', g'(U_X \cap U_{Y'})) \in \Xi_u; gU_X = g'U_X\}$ is an affine space. Fix $Y' \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}$ such that $\text{pos}(X, Y') = u$. (The set of such Y' is a homogeneous space $U_X/(U_X \cap Y)$, hence is an affine space of dimension δ .) It is enough to show that

$\{g'(U_X \cap U_{Y'}) \in (N_D X \cap N_D Y)/(U_X \cap U_Y); g' \in gU_X\}$ is a point.

Now $g = g_0 v_0$ where $g_0 \in N_D X \cap N_D Y$, $v_0 \in U_X$. It is enough to show that $\{v \in U_X; v_0 v \in N_G X \cap N_G Y\}/(U_X \cap U_Y)$ is a point or that $(U_X \cap N_G X \cap N_G Y)/(U_X \cap U_Y)$ is a point or that $U_X \cap Y = U_X \cap U_Y$. This is clear.

38.7. Let $u \in {}^K\mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon$. We set $H = K \cap \text{Ad}(u)K'$, $H' = K' \cap \text{Ad}(u^{-1})K = \text{Ad}(u^{-1})H$. Let $\Psi'_u = \Psi_u[[\delta]] : \mathcal{D}(V_{H,D}) \rightarrow \mathcal{D}(V_{H',D})$. We show:

(a) If $A \in CS(V_{H,D})$ then $\Psi'_u(A) \in CS(V_{H',D})$.

We have a commutative diagram in which the upper squares are cartesian and the left and right vertical arrows are smooth with connected fibres:

$$\begin{array}{ccccc}
 V_{H,D} & \xleftarrow{j} & \Xi_u & \xrightarrow{h} & V_{H',D} \\
 f'_3 \uparrow & & f'_4 \uparrow & & f'_5 \uparrow \\
 V'_H & \xleftarrow{j'} & \Xi''_u & \xrightarrow{h'} & V'_{H'} \\
 f_3 \downarrow & & f_4 \downarrow & & f_5 \downarrow \\
 V_H^u & \xleftarrow{j^u} & \Xi'_u & \xrightarrow{h^u} & V_{H'}^u
 \end{array}$$

Here

$$\begin{aligned}
 V_H^u &= \{(X, gU_X) \in V_{H,D}; X = X_u\}, \quad V_{H'}^u = \{(Y, gU_Y) \in V_{H',D}; Y = Y_u\}, \\
 \Xi'_u &= \{(X, Y, g(U_X \cap U_Y)) \in \Xi_u; X = X_u, Y = Y_u\}, \\
 V'_H &= G^0/U_{X_u} \times V_H^u, \quad V'_{H'} = G^0/U_{Y_u} \times V_{H'}^u, \\
 \Xi''_u &= G^0/(U_{X_u} \cap U_{Y_u}) \times \Xi'_u,
 \end{aligned}$$

$$\begin{aligned}
& j^u, h^u \text{ are the restrictions of } j, h, \\
& j'(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) = (xU_{X_u}, (X_u, gU_{X_u})), \\
& h'(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) = (xU_{Y_u}, (Y_u, gU_{Y_u})), \\
& f'_3(xU_{X_u}, (X_u, gU_{X_u})) = (xX_ux^{-1}, xgx^{-1}U_{xX_ux^{-1}}), \\
& f_3(xU_{X_u}, (X_u, gU_{X_u})) = (X_u, gU_{X_u}), \\
& f'_4(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) = (xX_ux^{-1}, xY_ux^{-1}, xgx^{-1}(U_{xX_ux^{-1}} \cap \\
& U_{xY_ux^{-1}})), \\
& f'_5(xU_{Y_u}, (Y_u, gU_{Y_u})) = (xY_ux^{-1}, xgx^{-1}U_{xY_ux^{-1}}), \\
& f_4(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) = (X_u, Y_u, g(U_{X_u} \cap U_{Y_u})), \\
& f_5(xY_u, (Y_u, gU_{Y_u})) = (Y_u, gU_{Y_u}).
\end{aligned}$$

We may identify $V_H^u = N_D X_u / U_{X_u}$, $V_{H'}^u = N_D Y_u / U_{Y_u}$ in an obvious way. We can find $C \in CS(N_D X_u / U_{X_u})$ such that $f_3'^\star A = f_3^\star C$. Hence $f_3'^\star A[\dim X_u / U_{X_u}] = f_3^\star C[\dim G / U_{X_u}]$ and $f_3'^\star A = f_3^\star A_1$ where $A_1 = C[\alpha_u]$. Let $A_2 = h_!^u j^{u*} A_1$, $A_2' = h_! j^* A$. Using 38.6(a) and the fact that h^u is an isomorphism, we have

$$\begin{aligned}
f_5'^\star A_2' &= f_5'^\star h_! j^* A = h_! f_4'^\star j^* A = h_! j'^\star f_3'^\star A = h_! j'^\star f_3^\star A_1 = h_! f_4'^\star j^{u*} A_1 \\
&= h_! f_4'^\star h^{u*} h_!^u j^{u*} A_1 = h_! h'^\star f_5^* h_!^u j^{u*} A_1 = f_5^* h_!^u j^{u*} A_1[[-\delta]] = f_5^* A_2[[-\delta]].
\end{aligned}$$

Thus, $f_5^* A_2[[-\delta]] = f_5'^\star A_2'$. Hence

$$f_5^\star A_2[-\dim G^0 / U_{Y_u}][[-\delta]] = f_5'^\star A_2'[-\dim Y_u / U_{Y_u}], f_5^\star A_2[-\alpha_u] = f_5'^\star A_2'[[\delta]].$$

Since h^u, j^u are isomorphisms we see that $A_2[-\alpha_u]$ is a perverse sheaf. Hence so is $f_5^\star A_2[-\alpha_u]$. Hence $f_5'^\star A_2'[[\delta]]$ is perverse. Hence $A_2'[[\delta]]$ is perverse and $h_! j^* A[[\delta]]$ is perverse. To show that $h_! j^* A[[\delta]] = A_2'[[\delta]] \in CS(V_{H',D})$, it is enough to show that $A_2[-\alpha_u] \in CS(N_D Y_u / U_{Y_u})$ or that $h_!^u j^{u*} C \in CS(N_D Y_u / U_{Y_u})$.

Let M be a common Levi subgroup of X_u, Y_u . Let $\tilde{M} = N_G X_u \cap N_G Y_u \cap N_G M$. Then $\tilde{M}$ is a reductive group with $\tilde{M}^0 = M$ and $\tilde{M}^1 = N_D X_u \cap N_D Y_u \cap N_D M$ is a connected component of $\tilde{M}$. Moreover, the obvious maps $\tilde{M}^1 \rightarrow N_D X_u / U_{X_u}$, $\tilde{M}^1 \rightarrow N_D Y_u / U_{Y_u}$, $\tilde{M}^1 \rightarrow (N_D X_u \cap N_D Y_u) / (U_{X_u} \cap U_{Y_u})$ are isomorphisms (see 1.25). Hence the bottom row $V_H^u \xleftarrow{j^u} \Xi'_u \xrightarrow{h^u} V_{H'}^u$ of the commutative diagram above may be identified with $\tilde{M}^1 \leftarrow \tilde{M}^1 \rightarrow \tilde{M}^1$ where both maps are the identity. Thus $h_!^u j^{u*} C \in CS(N_D Y_u / U_{Y_u})$ follows immediately from $C \in CS(N_D X_u / U_{X_u})$. This proves (a).

We now show:

(b) For A as above we have $f_{H',J} \Psi'_u(A) = f_{H,J}(A)$.

Let $Q \in \mathcal{P}_J$ be such that $X_u \subset Q \supset Y_u$. Let M_1 be the unique Levi of Q such that $M \subset M_1$. Let $\tilde{M}_1 = N_G Q \cap N_G M_1$. Then $\tilde{M}_1$ is a reductive group with $\tilde{M}_1^0 = M_1$ and $\tilde{M}_1^1 = N_D Q \cap N_D M_1$ is a connected component of $\tilde{M}_1$. Moreover, $X_u \cap \tilde{M}_1, Y_u \cap \tilde{M}_1$ are parabolic subgroups of M_1 with a common Levi, M . Let $C \in CS(N_D X_u / U_{X_u}) = CS(\tilde{M}_1^1)$ be as above. We may assume that A is simple so that C is also simple. Using the proof of (a) and that of 38.2 we see that it

is enough to verify that $\text{ind}_{\tilde{M}^1}^{\tilde{M}^1}(C)$ (defined in terms of the parabolic $X_u \cap \tilde{M}_1$) is isomorphic to $\text{ind}_{\tilde{M}^1}^{\tilde{M}^1}(C)$ (defined in terms of the parabolic $Y_u \cap \tilde{M}_1$). Since C is an admissible complex on $\tilde{M}^1$ (see 30.12), this follows from 27.2(d) which shows (for G instead of $\tilde{M}_1$) that $\text{ind}_{\tilde{M}^1}^{\tilde{M}^1}(C)$ can be defined without reference to a choice of parabolic. This proves (b).

Proposition 38.8. *Let K, K', J be as in 38.4. Let $A \in CS(V_{K,D})$. We have*

$$e_{K',J} f_{K,J} A \cong \bigoplus_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} f_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi'_u e_{K \cap \text{Ad}(u)K', K} A$$

in $CS(V_{K',D})$.

We set $\mathfrak{C} = e_{K',J} f_{K,J} A \in \mathcal{D}(V_{K,D})$. For any $u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon$ we set $\mathfrak{C}_u = f_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi'_u e_{K \cap \text{Ad}(u)K', K} A \in \mathcal{D}(V_{K',D})$. Assume that we can show that

$$\mathfrak{C} \simeq \{\mathfrak{C}_u; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}.$$

From the definition of $\simeq$ (see 32.15) it would then follow that

$$\sum_i (-1)^i ({}^p H^i(\mathfrak{C})) = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} \sum_i (-1)^i ({}^p H^i(\mathfrak{C}_u))$$

in the Grothendieck group of the category of perverse sheaves on $V_{K',D}$. From 38.2(a), 38.3(a), 38.7(a) we see that $\mathfrak{C}, \mathfrak{C}_u \in CS(V_{K',D})$ (hence are perverse sheaves); hence the previous equality implies that $\mathfrak{C} = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} \mathfrak{C}_u$ in the Grothendieck group of the category of perverse sheaves on $V_{K',D}$. Since $\mathfrak{C}, \mathfrak{C}_u$ are semisimple perverse sheaves (being in $CS(V_{K',D})$) it follows that $\mathfrak{C} \cong \bigoplus_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} \mathfrak{C}_u$, as desired. ■

Assume that $P \in \mathcal{P}_K$ contains X_u and $R \in \mathcal{P}_{K'}$ contains Y_u so that $\text{pos}(P, R) = u$. (X_u, Y_u as in 38.6.) Let $Q \in \mathcal{P}_J$ be such that $P \subset Q \supset R$. Let $\beta = \dim U_Q$. We have

$$\mathfrak{C} = \tilde{\mathfrak{C}}[\alpha + \alpha' - 2\beta](\alpha' - \beta),$$

$$\mathfrak{C}_u = \tilde{\mathfrak{C}}_u\alpha_u - \alpha[\alpha_u - \alpha'] [2\delta](\delta).$$

(Notation of 38.5.) Hence it is enough to show

$$\tilde{\mathfrak{C}}[\alpha + \alpha' - 2\beta](\alpha' - \beta) \simeq \{\tilde{\mathfrak{C}}_u[2\alpha_u - \alpha - \alpha' + 2\delta](\alpha_u - \alpha + \delta); u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}$$

or that

$$\tilde{\mathfrak{C}} \simeq \{\tilde{\mathfrak{C}}_u[2\alpha_u - 2\alpha - 2\alpha' + 2\delta - 2\beta](\alpha_u - \alpha - \alpha' + \delta + \beta); u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}.$$

By 38.5, it is enough to show that for any u we have $\alpha_u - \alpha - \alpha' + \delta + \beta = -m_u$ or that

$$\dim U_P + \dim U_R - \dim(U_P \cap U_R) = \dim U_{X_u} + \dim U_{Y_u} - \dim(U_{X_u} \cap U_{Y_u}).$$

It is enough to show that

$$\begin{aligned} & \dim \operatorname{Lie} U_P + \dim \operatorname{Lie} U_R - \dim(\operatorname{Lie} U_P \cap \operatorname{Lie} U_R) \\ &= \dim \operatorname{Lie} U_{X_u} + \dim \operatorname{Lie} U_{Y_u} - \dim(\operatorname{Lie} U_{X_u} \cap \operatorname{Lie} U_{Y_u}) \end{aligned}$$

or that $\dim(\operatorname{Lie} U_P + \operatorname{Lie} U_R) = \dim(\operatorname{Lie} U_{X_u} + \operatorname{Lie} U_{Y_u})$. We have

$$\operatorname{Lie} U_{X_u} = \operatorname{Lie} U_P + (\operatorname{Lie} P \cap \operatorname{Lie} U_R), \operatorname{Lie} U_{Y_u} = \operatorname{Lie} U_R + (\operatorname{Lie} R \cap \operatorname{Lie} U_P)$$

hence $\operatorname{Lie} U_{X_u} + \operatorname{Lie} U_{Y_u} \subset \operatorname{Lie} U_P + \operatorname{Lie} U_R$. The opposite inclusion is clear since $\operatorname{Lie} U_P \subset \operatorname{Lie} U_{X_u}, \operatorname{Lie} U_R \subset \operatorname{Lie} U_{Y_u}$. Thus we have

$$\operatorname{Lie} U_{X_u} + \operatorname{Lie} U_{Y_u} = \operatorname{Lie} U_P + \operatorname{Lie} U_R.$$

This completes the proof.

38.9. For $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ let $\mathcal{K}(V_{J,D})$ be the Grothendieck group of $CS(V_{J,D})$. Similarly let $\mathcal{K}(D)$ be the Grothendieck group of $CS(D)$. For $A, A' \in CS(V_{J,D})$ we set $(A, A') = \dim \operatorname{Hom}(A, A')$. This induces a symmetric bilinear pairing $(\cdot, \cdot)_J : \mathcal{K}(V_{J,D}) \times \mathcal{K}(V_{J,D}) \rightarrow \mathbf{Z}$.

For $J \subset J' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J'$, the functors $f_{J,J'} : CS(V_{J,D}) \rightarrow CS(V_{J',D})$ and $e_{J,J'} : CS(V_{J',D}) \rightarrow CS(V_{J,D})$ are compatible with direct sums hence they induce homomorphisms $\mathcal{K}(V_{J,D}) \rightarrow \mathcal{K}(V_{J',D}), \mathcal{K}(V_{J',D}) \rightarrow \mathcal{K}(V_{J,D})$ denoted again by $f_{J,J'}, e_{J,J'}$. From 30.5 we see that

$$(a) \quad (e_{J,J'} A', A)_J = (A', f_{J,J'} A)_{J'}$$

for $A \in \mathcal{K}(V_{J,D}), A' \in \mathcal{K}(V_{J',D})$.

In the setup of 38.7, for $u \in {}^K \mathbf{W}^{K'} \in \mathbf{W}_J^\epsilon$, the functor

$$\Psi'_u : CS(V_{K \cap \operatorname{Ad}(u)K', D}) \rightarrow CS(V_{K' \cap \operatorname{Ad}(u^{-1})K, D})$$

is compatible with direct sums hence induces a homomorphism

$$\mathcal{K}(V_{K \cap \operatorname{Ad}(u)K', D}) \rightarrow \mathcal{K}(V_{K' \cap \operatorname{Ad}(u^{-1})K, D})$$

denoted again by Ψ'_u . Below we shall need the following identity:

$$(b) \quad \sum_{\substack{K'; K' \subset J \\ \epsilon(K')=K'}} \sum_{\substack{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon \\ K \cap \operatorname{Ad}(u)K'=H}} (-1)^{|K'_\epsilon|} = (-1)^{|H_\epsilon|}$$

for any $H \subset K \subset J \subset \mathbf{I}$ such that $\epsilon(H) = h, \epsilon(K) = K, \epsilon(J) = J$. In the case where $J = \mathbf{I}, \epsilon = 1$ this is proved in [Cu, 2.5]; the general case can be reduced to this special case by replacing $\mathbf{W}$ by $\mathbf{W}_J^\epsilon$ which is itself a Weyl group with simple reflections in bijection with J_ϵ .

38.10. For $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ we define a homomorphism $\mathbf{d}_J : \mathcal{K}(V_{J,D}) \rightarrow \mathcal{K}(V_{J,D})$ by

$$(a) \quad \mathbf{d}_J A = \sum_{K; K \subset J, \epsilon(K)=K} (-1)^{|K_\epsilon|} f_{K,J} e_{K,J} A.$$

Using 38.9(a) we see that for $A, A' \in \mathcal{K}(V_{J,D})$ we have

$$(b) \quad (\mathbf{d}_J(A), A')_J = (A, \mathbf{d}_J(A'))_J.$$

We show that, for $K \subset J \subset \mathbf{I}$ such that $\epsilon(K) = K$, $\epsilon(J) = J$ and $A \in \mathcal{K}(V_{K,D})$, we have

$$(c) \quad \mathbf{d}_J f_{K,J} A = f_{K,J} \mathbf{d}_K A.$$

Using 38.8, 38.7(b), 38.9(b), 38.1(a) we have

$$\begin{aligned}
\mathbf{d}_J f_{K,J} A &= \sum_{K'; K' \subset J, \epsilon(K')=K'} (-1)^{|K'_\epsilon|} f_{K',J} e_{K',J} f_{K,J} A \\
&= \sum_{\substack{K'; K' \subset J \\ \epsilon(K')=K' \\ u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon}} (-1)^{|K'_\epsilon|} f_{K',J} f_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi'_u e_{K \cap \text{Ad}(u)K', K} A \\
&= \sum_{\substack{H; H \subset J, \epsilon(H)=H \\ K' \\ u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon \\ K' \subset J, \epsilon(K')=K' \\ K \cap \text{Ad}(u)K'=H}} (-1)^{|K'_\epsilon|} f_{\text{Ad}(u^{-1})H, J} \Psi'_u e_{H, K} A \\
&= \sum_{\substack{H; H \subset J, \epsilon(H)=H \\ K'; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon \\ K' \subset J, \epsilon(K')=K' \\ K \cap \text{Ad}(u)K'=H}} (-1)^{|K'_\epsilon|} f_{H, J} e_{H, K} A \\
&= \sum_{\substack{H; H \subset J, \epsilon(H)=H \\ K'; K' \subset J, \epsilon(K')=K' \\ u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon \\ K \cap \text{Ad}(u)K'=H}} (-1)^{|K'_\epsilon|} f_{H, J} e_{H, K} A = \sum_{H; H \subset K, \epsilon(H)=H} (-1)^{|H_\epsilon|} f_{H, J} e_{H, K} A \\
&= \sum_{H; H \subset K, \epsilon(H)=H} (-1)^{|H_\epsilon|} f_{K, J} f_{H, K} e_{H, K} A = f_{K, J} \mathbf{d}_K A
\end{aligned}$$

and (c) is proved.

We show that, for $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ and $A \in \mathcal{K}(V_{J,D})$, we have

$$(d) \quad \mathbf{d}_J \mathbf{d}_J A = A.$$

Using (c), 38.1(a), we have

$$\begin{aligned}
\mathbf{d}_J \mathbf{d}_J A &= \sum_{K; K \subset J, \epsilon(K)=K} (-1)^{|K_\epsilon|} \mathbf{d}_J f_{K,J} e_{K,J} A \\
&= \sum_{K; K \subset J, \epsilon(K)=K} (-1)^{|K_\epsilon|} f_{K,J} \mathbf{d}_K e_{K,J} A \\
&= \sum_{K, K'; K' \subset K \subset J, \epsilon(K)=K, \epsilon(K')=K'} (-1)^{|K_\epsilon|} (-1)^{|K'_\epsilon|} f_{K,J} f_{K',K} e_{K',K} e_{K,J} A \\
&= \sum_{K, K'; K' \subset K \subset J, \epsilon(K)=K, \epsilon(K')=K'} (-1)^{|K_\epsilon|} (-1)^{|K'_\epsilon|} f_{K',J} e_{K',J} A \\
&= \sum_{K'; K' \subset J, \epsilon(K')=K'} (-1)^{|K'_\epsilon|} \sum_{K; K' \subset K \subset J, \epsilon(K)=K} (-1)^{|K_\epsilon|} f_{K',J} e_{K',J} A \\
&= \sum_{K'; K' \subset J, \epsilon(K')=K'} (-1)^{|K'_\epsilon|} \delta_{K',J} (-1)^{|K'_\epsilon|} f_{K',J} e_{K',J} A = f_{J,J} e_{J,J} A = A
\end{aligned}$$

and (d) is proved.

We show that, for $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ and $A, A' \in \mathcal{K}(V_{J,D})$, we have

$$(e) \quad (\mathbf{d}_J A, \mathbf{d}_J A')_J = (A, A')_J.$$

Using (b),(d) we have $(\mathbf{d}_J A, \mathbf{d}_J A')_J = (A, \mathbf{d}_J \mathbf{d}_J A')_J = (A, A')_J$ as desired.

38.11. We write $\mathbf{d}$ instead of $\mathbf{d}_{\mathbf{I}}$ and $(,)$ instead of $(,)_{\mathbf{I}}$. We call $\mathbf{d}$ the duality operator on character sheaves. If A is a character sheaf on $D = V_{\mathbf{I},D}$, then $(A, A) = 1$ (where A is regarded as an element of $\mathcal{K}(D)$) hence, by 38.10(e), we have $(\mathbf{d}(A), \mathbf{d}(A)) = 1$. Since $\mathbf{d}(A)$ is a $\mathbf{Z}$ -linear combination of isomorphism classes of character sheaves (which form an orthonormal basis of $\mathcal{K}(D)$ for $(,)$) it follows that $\mathbf{d}(A) = \pm A'$ where A' is a well defined character sheaf on D (up to isomorphism). The sign can be described as follows. By 30.6(d) we can find a parabolic P_0 of G^0 such that $N_D P_0 \neq \emptyset$ and a cuspidal character sheaf A_0 on $D_0 := N_D P_0 / U_{P_0}$ such that A is a direct summand of $\text{ind}_{D_0}^D(A_0)$. We have $P_0 \in \mathcal{P}_J$ where $J \subset \mathbf{I}$, $\epsilon(J) = J$. Then

$$(a) \quad \mathbf{d}(A) = (-1)^{|J_\epsilon|} A'.$$

Indeed let $A_0^b \in CS(V_{D,J})$ be the perverse sheaf corresponding to A_0 as in 30.3. Then $e_{J',J} A_0^b = 0$ for any $J' \subsetneq J$ such that $\epsilon(J') = J'$, see 38.2. Hence $\mathbf{d}_J(A_0^b) = (-1)^{|J_\epsilon|} A_0^b$. Now A is a direct summand of $f_{J,\mathbf{I}} A_0^b$ and by 38.10(c) we have

$$\mathbf{d} f_{J,\mathbf{I}} A_0^b = f_{J,\mathbf{I}} \mathbf{d}_J A_0^b = (-1)^{|J_\epsilon|} f_{J,\mathbf{I}} A_0^b.$$

In $\mathcal{K}(D)$ we have $f_{J,\mathbf{I}} A_0^b = \sum_{k=1}^{k_0} n_k A_k$ where A_k are distinct character sheaves on D , $n_k \in \mathbf{Z}_{>0}$ and $A_1 = A$. We have $\mathbf{d}(A_k) = \iota_k A'_k$ where $\iota_k = \pm 1$ and

A'_k are distinct character sheaves on D , $n_k \in \mathbf{Z}_{>0}$ and $A_1 = A$. We see that $\sum_k n_k \iota_k A'_k = (-1)^{|J_\epsilon|} \sum_k n_k A_k$. Since $\{A_k\}$ and $\{A'_k\}$ are parts of the same basis of $\mathcal{K}(D)$ we see that $\iota_k / (-1)^{|J_\epsilon|} > 0$ for any k . Hence $\iota_k = (-1)^{|J_\epsilon|}$ for any k . In particular this holds for $k = 1$ and (a) follows.

Note that, by 38.10(d), $A \mapsto A'$ is an involution of the set of isomorphism classes of character sheaves on D .

38.12. The definition of the duality operator for character sheaves in 38.11 is entirely similar to that of a duality operator for representations of a reductive group over a finite field (given again by an alternating sum of compositions of a parabolic restriction and a parabolic induction) which was found by the author in 1977, who conjectured that it takes irreducibles to $\pm$ irreducibles and is involutive. In 1977 I communicated this conjecture to C.W.Curtis and N.Kawanaka, see [Ka, p.412]; the conjecture was proved in [Cu],[Al] and in [Ka].

39. QUASI-RATIONALITY

39.1. The main result of this section is Proposition 39.7 which gives a quasi-rationality property of representations of certain extensions of a Weyl group. This is needed to prove a key property of character sheaves (Corollary 39.8).

Let W, I be a Weyl group (I is the set of simple reflections). We have canonically $W = \prod_{j \in J} W_j$ where W_j is an irreducible Weyl group (with set of simple reflections $I_j = I \cap W_j$). We identify W_j with a subgroup of W . Let $R(W) = \cup_{w \in W} w I w^{-1}$. We have $R(W) = \sqcup_{j \in J} R(W_j)$. For any $j \in J$ the set $R(W_j)$ is a single W_j -conjugacy class if W_j is of type A, D or E ; it is a union of two W_j -conjugacy classes, otherwise. A subset $\mathcal{X}$ of $R(W)$ is said to be *special* if $\mathcal{X} = \sqcup_{j \in J} \mathcal{X}_j$ where $\mathcal{X}_j$ is a W_j -conjugacy class in $R(W_j)$. Clearly, a special subset of $R(W)$ exists; we assume that a special subset $\mathcal{X}$ of $R(W)$ is given. Let $\text{Aut}(W, I, \mathcal{X})$ be the group of automorphisms of W that preserve I and $\mathcal{X}$.

Lemma 39.2. *Let Γ be a finite group. Let $\gamma \mapsto \rho_\gamma$ be a homomorphism $\Gamma \rightarrow \text{Aut}(W, I, \mathcal{X})$. Let E be a simple $\mathbf{Q}[W]$ -module such that $\text{tr}(\rho_\gamma(w), E) = \text{tr}(w, E)$ for any $\gamma \in \Gamma, w \in W$. Assume that either*

(i) $|J| = 1$, or

(ii) Γ is an extension of a cyclic group by a cyclic group.

Then there exists a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$, $\gamma \mapsto t_\gamma$ such that $t_\gamma(w(e)) = \rho_\gamma(w)t_\gamma(e)$ for any $\gamma \in \Gamma, w \in W, e \in E$.

Γ acts on J by $\gamma : j \mapsto \gamma(j)$ where $W_{\gamma(j)} = \rho_\gamma(W_j)$. We may identify $E = \otimes_{j \in J} E_j$ as $\mathbf{Q}[W]$ -modules where E_j is a simple $\mathbf{Q}[W_j]$ -module for any $j \in J$. From our assumption we see that for any $j \in J, \gamma \in \Gamma$ there exists a $\mathbf{Q}$ -linear isomorphism

(a) $H_j^\gamma : E_j \xrightarrow{\sim} E_{\gamma(j)}$ with $H_j^\gamma(w_j e) = \rho_\gamma(w_j) H_j^\gamma(e)$ for all $e \in E_j, w_j \in W_j$; moreover, H_j^γ is unique up to multiplication by an element of $\mathbf{Q}^*$. It follows that

(b) $H_{\gamma(j)}^{\gamma'} H_j^\gamma \in \mathbf{Q}^* H_j^{\gamma'\gamma}$ for $j \in J$ and $\gamma, \gamma' \in \Gamma$.

For $\gamma \in \Gamma$ we define $\tilde{t}_\gamma \in \text{Aut}_{\mathbf{Q}}(E)$ by

(c) $\tilde{t}_\gamma(\otimes_j e_j) = \otimes_j e'_j$ where $e_j \in E_j$ and $e'_j = H_{\gamma^{-1}(j)}^\gamma(e_{\gamma^{-1}(j)}) \in E_j$.

From definitions we have $\tilde{t}_\gamma(w(e)) = \rho_\gamma(w)\tilde{t}_\gamma(e)$ for any $w \in W, e \in E$; moreover for $\gamma, \gamma' \in \Gamma$, the maps $\tilde{t}_{\gamma'\gamma}, \tilde{t}_{\gamma'}\tilde{t}_\gamma$ are equal up to a factor in $\mathbf{Q}^*$. Thus the maps $\tilde{t}_\gamma$ provide a homomorphism $\Gamma \rightarrow PGL(E)$ rather than a homomorphism $\Gamma \rightarrow GL(E)$.

We prove the lemma in the setup of (i). Replacing Γ by its image under $\gamma \mapsto \rho_\gamma$ we may assume that $\Gamma \subset \text{Aut}(W, I, \mathcal{X})$ and $\gamma \mapsto \rho_\gamma$ is the inclusion. We form the semidirect product $W\Gamma$ with W normal. It is enough to show that E extends to a $W\Gamma$ -module. We may assume that $\Gamma \neq \{1\}$. Then W is of type $A_n (n \geq 2), D_n$ or E_6 . If $\Gamma = \text{Aut}(W, I, \mathcal{X})$ then $W\Gamma$ is itself a Weyl group, of type $A_n \times A_1, D_{2n+1} \times A_1, B_{2n}, F_4, E_6 \times A_1$ for W of type $A_n (n \geq 2), D_{2n+1} (n \geq 2), D_{2n} (n \geq 3), D_4, E_6$ respectively and the desired result follows easily from the known properties of representations of such Weyl groups (in particular, from their rationality); the same applies if W is of type D_4 and $|\Gamma| = 2$ (in this case, $W\Gamma$ is a Weyl group of type B_4). In the only remaining case (W of type $D_4, |\Gamma| = 3$), the result follows by an argument in [L14, 3.2].

Next we prove the lemma in the setup of (ii). Now Γ has two generators a, c and relations

$$a^M = 1, c^N = a^u, cac^{-1} = a^k$$

where M, N, k are integers ≥ 1 such that $k^N = 1 \pmod{M}$ and $u \in \mathbf{N}$ satisfies $uk = k \pmod{M}$.

We consider separately 3 cases in increasing order of generality.

Case 1. Assume that J is a single orbit of $a : J \rightarrow J$. We may identify $J = \mathbf{Z}/m$ so that $a(i) = i+1, c(i) = ki - r$ for $i \in \mathbf{Z}/m$; here $r \in \mathbf{N}$ is independent of i . Since $a^M = 1 : J \rightarrow J$ we see that m divides M . For any i we have $i+u = c^N(i) = k^N i - (1+k+k^2+\dots+k^{N-1})r$ in $\mathbf{Z}/m$. Since $k^N = 1 \pmod{M}$ (hence $k^N = 1 \pmod{m}$) we have $u + (1+k+k^2+\dots+k^{N-1})r = fm$ for some $f \in \mathbf{N}$. We set $c' = a^r c$. Then $c'(i) = ki$ for $i \in \mathbf{Z}/m$. Let Γ' be the subgroup of Γ generated by a^m, c' . In Γ' we have

$$(a^m)^{M/m} = 1, c'^N = (a^m)^f, c' a^m c'^{-1} = (a^m)^k.$$

Since $a^m(0) = 0, c'(0) = 0$, the action of Γ on W restricts to an action of Γ' on W_0 . Using the lemma (setup of (i)) for W_0, E_0, Γ' instead of W, E, Γ we obtain a homomorphism $\Gamma' \rightarrow \text{Aut}_{\mathbf{Q}}(E_0)$ such that, denoting by s_0, s_1 the images of a^m, c' under this homomorphism, we have $s_0^{M/m} = 1, s_1^N = s_0^f, s_1 s_0 s_1^{-1} = s_0^k, s_0(w(e)) = \rho_a^m(w)s_0(e), s_1(w(e)) = \rho_{c'}(w)s_1(e)$ for any $w \in W_0, e \in E_0$.

For $i \in \mathbf{Z}/m$, let $H_i^a : E_i \xrightarrow{\sim} E_{i+1}$ be as in (a). Then $s_2 := H_{m-1}^a \dots H_1^a H_0^a : E_0 \xrightarrow{\sim} E_0$ satisfies $s_2(w(e)) = \rho_a^m(w)s_2(e)$ for any $w \in W_0, e \in E_0$. Hence $s_2^{-1}s_0(w(e)) = w(s_2^{-1}s_0(e))$ for any $w \in W_0, e \in E_0$. By the absolute irreducibility of the W_0 -module E_0 we see that $s_2^{-1}s_0 : E_0 \rightarrow E_0$ is a $\mathbf{Q}^*$ -multiple of the identity map. Hence, replacing H_0^a by a $\mathbf{Q}^*$ -multiple, we may assume that $s_2 = s_0$ that is,

$H_{m-1}^a \dots H_1^a H_0^a = s_0$. For any $h \in \mathbf{N}$ we set

$b_h = H_{hk-1}^a H_{hk-2}^a \dots H_1^a H_0^a s_1 (H_{h-1}^a \dots H_1^a H_0^a)^{-1} : E_h \xrightarrow{\sim} E_{kh}$
(there are hk factors to the left of s_1 and h factors to the right of s_1). For $h \geq 1$ we have

$$(d) \quad b_h = H_{hk-1}^a H_{hk-2}^a \dots H_{hk-k}^a b_{h-1} (H_{h-1}^a)^{-1}.$$

We show that $b_{h+m} = b_h$ for $h \in \mathbf{N}$. We argue by induction on h . Assume first that $h = 0$. We must show that $H_{mk-1}^a H_{mk-2}^a \dots H_1^a H_0^a s_1 (H_{m-1}^a \dots H_1^a H_0^a)^{-1} = s_1$; this is a reformulation of the already known equality $s_0^k s_1 s_0^{-1} = s_1$. Assume next that $h \geq 1$. Using (d) twice and the induction hypothesis we have

$$\begin{aligned} b_{h+m} &= H_{hk+m-1}^a H_{hk+m-2}^a \dots H_{hk+m-k}^a b_{h+m-1} (H_{h+m-1}^a)^{-1} \\ &= H_{hk-1}^a H_{hk-2}^a \dots H_{hk-k}^a b_{h-1} (H_{h-1}^a)^{-1} = b_h, \end{aligned}$$

as desired. We see that b_h depends only on the image of h in $\mathbf{Z}/m$.

We set $k = k^{N-1}$. Then $kk' = 1 \pmod{M}$. Define $t_a, t_{c'} \in \text{Aut}_{\mathbf{Q}}(E)$ by $t_a(\otimes_{i \in \mathbf{Z}/m} e_i) = \otimes_{i \in \mathbf{Z}/m} e'_i$ where $e_i \in E_i$ and $e'_i = H_{i-1}^a(e_{i-1}) \in E_i$,

$$t_{c'}(\otimes_{i \in \mathbf{Z}/m} e_i) = \otimes_{i \in \mathbf{Z}/m} e''_i \text{ where } e_i \in E_i \text{ and } e''_i = b_{ik'}(e_{ik'}) \in E_{ik'k} = E_i.$$

From definitions we have $t_a(w(e)) = \rho_a(w)t_a(e)$, $t_{c'}(w(e)) = \rho_{c'}(w)t_{c'}(e)$ for any $w \in W, e \in E$.

We have $t_{c'} t_a t_{c'}^{-1} = t_a^k$. This follows from the identity

$$b_{hk'} H_{hk'-1}^a b_{hk'-1}^{-1} = H_{h-1}^a H_{h-2}^a \dots H_{h-k}^a : E_{h-k} \rightarrow E_h$$

for $h = 0, 1, \dots, m-1$ (here b_{-1} is taken to be b_{m-1}); an equivalent identity is

$$b_i H_{i-1}^a b_{i-1}^{-1} = H_{ik-1}^a H_{ik-2}^a \dots H_{ik-k}^a : E_{ik-k} \rightarrow E_{ik}$$

for $i = 0, 1, \dots, m-1$, which is the same as (d).

We have $t_a^M = 1$. This follows from the identity $H_{i+M-1}^a \dots H_{i+1}^a H_i^a = 1 : E_i \rightarrow E_i$ for $i = 0, 1, \dots, m-1$ which is equivalent to the known equality $s_0^{M/m} = 1$.

We show that

$$(e) \quad t_{c'}^N = t_a^{mf}.$$

This follows from the identity

$$b_{ik'} b_{ik'^2} \dots b_{ik'^N} = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a \text{ for } i = 0, 1, \dots, m-1$$

or equivalently

$$b_{ik^{N-1}} b_{ik^{N-2}} \dots b_i = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a \text{ for } i = 0, 1, \dots, m-1.$$

This is the same as

$$H_{ik^N-1}^a H_{ik^N-2}^a \dots H_1^a H_0^a s_1^N (H_{i-1}^a \dots H_1^a H_0^a)^{-1} = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a$$

(with ik^N factors to the left of s_1^N). We have $ik^N - i = iMl$ for some $l \in \mathbf{N}$. From $s_0^{M/m} = 1$ we deduce $s_0^{iMl/m} = 1$ hence $H_{ik^N-1}^a H_{ik^N-2}^a \dots H_i^a = 1$ (with $ik^N - i$ factors). Since $s_1^N = s_0^f$, it remains to show

$$H_{i-1}^a \dots H_1^a H_0^a s_0^f (H_{i-1}^a \dots H_1^a H_0^a)^{-1} = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a.$$

If $f = 0$ this is obvious. Assume now that $f \geq 1$. Then $i - mf < 0$ and we see that it is enough to show

$$s_0^f (H_{i-1}^a \dots H_1^a H_0^a)^{-1} = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a$$

that is,

$$s_0^f = H_{-1}^a H_{-2}^a \cdots H_{i-mf}^a (H_{i-1}^a \cdots H_1^a H_0^a).$$

This can be rewritten in the form

$$s_0^f = H_{-1}^a H_{-2}^a \cdots H_{i-mf}^a H_{i-mf-1}^a \cdots H_{1-mf}^a H_{-mf}^a$$

which follows from $s_0 = H_{m-1}^a \cdots H_1^a H_0^a$. This proves (e).

We set $t_c = t_a^{-r} t_{c'} \in \text{Aut}_{\mathbf{Q}}(E)$. From definitions we have $t_c(w(e)) = \rho_c(w) t_c(e)$ for any $w \in W, e \in E$.

We have $t_c t_a t_c^{-1} = t_a^k$. This follows from the identity $t_{c'} t_a t_{c'}^{-1} = t_a^k$.

We have $t_c^N = t_a^u$. Indeed,

$$(t_a^{-r} t_{c'})^N = t_a^{-r(1+k+k^2+\cdots+k^{N-1})} t_{c'}^N = t_a^{-mf+u} t_a^{mf} = t_a^u.$$

We see that $t_a, t_c \in \text{Aut}_{\mathbf{Q}}(E)$ satisfy the relations of Γ hence they define a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$. This has the required properties.

Case 2. Assume that J is a single Γ -orbit. Let $\langle a \rangle$ be the subgroup of Γ generated by a . If X is an $\langle a \rangle$ -orbit in J then cX is again an $\langle a \rangle$ -orbit. (We must show that, if $j \in X$ and $i \geq 1$, then $ca^i j, cj$ are in the same $\langle a \rangle$ -orbit. But $ca^i j = a^{ik} cj$.) Hence $c^h X$ is an $\langle a \rangle$ -orbit in J for any $h \in \mathbf{N}$. We can find an integer $z \geq 1$ such that $X, cX, \dots, c^{z-1}X$ are distinct and $c^z X = X$. (Clearly, z is a divisor of N .) Hence the notation $X_h = c^h X$ for $h \in \mathbf{Z}/z$ is meaningful. Now $X_0, X_1, \dots, X_{z-1}$ are distinct. The union $\cup_{h \in [0, z-1]} X_h$ is c -stable and a -stable hence it is equal to J (by our assumption on J). We see that $|J| = |X|z$. For $h \in \mathbf{Z}/z$ we set $W^h = \prod_{j \in X_h} W_j$, $E^h = \otimes_{j \in X_h} E_j$. We have naturally $W = \prod_{h \in [0, z-1]} W^h$, $E = \otimes_{h \in [0, z-1]} E^h$. Let Γ'' be the subgroup of Γ generated by a, c^z . In Γ'' we have

$$a^M = 1, (c^z)^{N/z} = a^u, c^z a c^{-z} = a^{k^z}.$$

Since $a(X_0) = X_0, c^z(X_0) = X_0$, the action of Γ on W restricts to an action of Γ'' on W^0 . Using case 1 for W^0, E^0, Γ'' instead of W, E, Γ we obtain a homomorphism $\Gamma'' \rightarrow \text{Aut}_{\mathbf{Q}}(E^0)$ such that, denoting by S_0, S_1 the images of a, c^z under this homomorphism, we have $S_0^M = 1, S_1^{N/z} = S_0^u, S_1 S_0 S_1^{-1} = S_0^{k^z}, S_0(w(e)) = \rho_a(w) S_0(e), S_1(w(e)) = \rho_c^z(w) S_1(e)$ for any $w \in W^0, e \in E^0$.

For $h \in \mathbf{Z}/z$ there exists a $\mathbf{Q}$ -linear isomorphism $K_h : E^h \xrightarrow{\sim} E^{h+1}$ with $K_h(we) = \rho_c(w) K_h(e)$ for all $e \in E^h, w \in W^h$ (note that $\rho_c(w) \in W^{h+1}$); moreover, K_h is unique up to multiplication by an element of $\mathbf{Q}^*$. (For example we can take K_h of the form $\otimes_{j \in X_h} H_j^c$ where $H_j^c : E_j \xrightarrow{\sim} E_{cj}$ are as in (a).) Then $S_2 := K_{z-1} \cdots K_1 K_0 : E^0 \xrightarrow{\sim} E^0$ satisfies $S_2(w(e)) = \rho_c^z(w) S_2(e)$ for any $w \in W^0, e \in E^0$. Hence $S_2^{-1} S_1(w(e)) = w(S_2^{-1} S_1(e))$ for any $w \in W^0, e \in E^0$. By the absolute irreducibility of the W^0 -module E^0 we see that $S_2^{-1} S_1 : E^0 \rightarrow E^0$ is a $\mathbf{Q}^*$ -multiple of the identity map. Hence, replacing K_0 by a $\mathbf{Q}^*$ -multiple, we may assume that $S_2 = S_1$ that is, $K_{z-1} \cdots K_1 K_0 = S_1$.

For any $h \in \mathbf{N}$ we define $\beta_{-h} \in \text{Aut}_{\mathbf{Q}}(E^{-h})$ by

$$\beta_{-h} = K_{-h}^{-1} \cdots K_{-2}^{-1} K_{-1}^{-1} S_0^{k^h} K_{-1} K_{-2} \cdots K_{-h}$$

(for $h = 0$ this is interpreted as $\beta_0 = S_0$). We show that $\beta_{-h-z} = \beta_{-h}$. We argue by induction on h . For $h = 0$ we must verify that

$$K_{-z}^{-1} \dots K_{-2}^{-1} K_{-1}^{-1} S_0^{k^z} K_{-1} K_{-2} \dots K_{-z} = S_0;$$

this follows from the known equality $S_1^{-1} S_0^{k^z} S_1 = S_0$. For $h \geq 1$ we have using the induction hypothesis

$$\beta_{-h-z} = K_{-h-z}^{-1} \beta_{-h-z+1}^k K_{-h-z} = K_{-h}^{-1} \beta_{-h+1}^k K_{-h} = \beta_{-z},$$

We see that β_{-h} depends only on the image of h in $\mathbf{Z}/z$. Define $t_a, t_c \in \text{Aut}_{\mathbf{Q}}(E)$ by

$$\tau_a(\otimes_{h \in \mathbf{Z}/z} e_h) = \otimes_{h \in \mathbf{Z}/z} e'_h \text{ where } e_h \in E^h \text{ and } e'_h = \beta_h(e_h) \in E^h,$$

$$\tau_c(\otimes_{h \in \mathbf{Z}/z} e_h) = \otimes_{h \in \mathbf{Z}/z} e''_h \text{ where } e_h \in E^h \text{ and } e''_h = K_{h-1}(e_{h-1}) \in E^h.$$

From definitions we have $t_a(w(e)) = \rho_a(w)t_a(e)$, $t_c(w(e)) = \rho_c(w)t_c(e)$ for any $w \in W, e \in E$.

We have $t_c t_a t_c^{-1} = t_a^k$. This follows from the identity $\beta_{h-1} = K_{h-1}^{-1} \beta_h^k K_{h-1}$ for $h \in \mathbf{Z}/z$.

We have $t_a^M = 1$. This follows from the identity $\beta_{-h}^M = 1$ for any $h \in \mathbf{N}$. An equivalent statement is $K_{-h}^{-1} \dots K_{-2}^{-1} K_{-1}^{-1} S_0^{k^h M} K_{-1} K_{-2} \dots K_{-h} = 1$ which follows from $S_0^M = 1$.

We show that $t_c^N = t_a^u$. It is enough to show that $K_{-h-1} K_{-h-2} \dots K_{-h-N} = b_{-h}^u : E_{-h} \rightarrow E_{-h}$ for $h = 0, 1, \dots, z-1$. An equivalent statement is

$$(K_{-h}^{-1} \dots K_{-1}^{-1})(K_{-1} K_{-2} \dots K_{-N})(K_{-1} \dots K_{-h}) = K_{-h}^{-1} \dots K_{-1}^{-1} S_0^{k^h u} K_{-1} \dots K_{-h}.$$

The left hand side is $(K_{-h}^{-1} \dots K_{-1}^{-1}) S_1^{N/z} (K_{-1} \dots K_{-h})$. It is enough to show that $S_1^{N/z} = S_0^{k^h u}$ or that $S_0^u = S_0^{k^h u}$ for $h = 0, 1, \dots, z-1$. Since $S_0^M = 1$ it is enough to show that $u = k^h u \pmod{M}$. This follows from $uk = u \pmod{M}$.

We see that $t_a, t_c \in \text{Aut}_{\mathbf{Q}}(E)$ satisfy the relations of Γ hence they define a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$. This has the required properties.

Case 3. We now consider the general case. For any Γ -orbit Y on J we set $W^Y = \prod_{j \in Y} W_j$, $E^Y = \otimes_{j \in Y} E_j$. We have naturally $W = \prod_Y W^Y$, $E = \otimes_Y E^Y$ where Y runs over the Γ -orbits in J . Now the Γ -action on W restricts to a Γ -action on W^Y for each Y . Using case 2 for W^Y, E^Y, Γ instead of W, E, Γ we obtain for each Y a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E^Y)$. We define a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$ by $\gamma : \otimes_Y e_Y \mapsto \otimes_Y (\gamma(e_Y))$; here $e_Y \in E^Y$. This has the required properties.

This completes the proof in the setup of (ii). The lemma is proved.

39.3. Let Γ be a finite group. Assume that Γ is a semidirect product of a normal subgroup Γ' with a cyclic group of order n with generator b . Let $\bar{\mathbf{U}}$ be an algebraic closed field of characteristic 0. Let E be a simple $\bar{\mathbf{U}}[\Gamma]$ -module which is isotypical as a $\bar{\mathbf{U}}[\Gamma']$ -module. We show that

(a) *E is simple as a $\bar{\mathbf{U}}[\Gamma']$ -module.*

There exists a simple $\bar{\mathbf{U}}[\Gamma']$ -module E' such that, setting $V = \text{Hom}_{\bar{\mathbf{U}}[\Gamma]}(E', E)$ we have $V \otimes E' \xrightarrow{\sim} E, f \otimes e' \mapsto f(e')$. For any $\gamma' \in \Gamma'$ we have $\text{tr}(b\gamma'b^{-1}, E') = \text{tr}(\gamma', E')$. Hence there exists $\xi \in \text{Aut}_{\bar{\mathbf{U}}}(E')$ such that $\xi(\gamma'e') = (b\gamma'b^{-1})\xi(e')$ for

$e' \in E', w' \in \Gamma'$. Then $\xi^n \in \text{Aut}_{\bar{\mathfrak{U}}}(E')$ commutes with the Γ' -action hence it is a scalar on E' . Replacing ξ by an $\bar{\mathfrak{U}}^*$ -multiple we can assume that $\xi^n = 1$. Then E' becomes a $\bar{\mathfrak{U}}[\Gamma]$ -module with b acting as ξ . Define $\eta : V \rightarrow V$ by $f \mapsto \eta(f)$ where $\eta(f)(e') = bf(\xi^{-1}(e'))$ for $e' \in E'$. We regard V as a $\bar{\mathfrak{U}}[\Gamma]$ -module in which Γ' acts trivially and b acts as η . Then the isomorphism $V \otimes E' \xrightarrow{\sim} E$ considered above is an isomorphism of $\bar{\mathfrak{U}}[\Gamma]$ -modules. Since E is simple it follows that V is a simple $\bar{\mathfrak{U}}[\Gamma]$ -module. Since Γ acts on V through a cyclic quotient, we see that $\dim V = 1$. It follows that $E \cong E'$ as a $\bar{\mathfrak{U}}[\Gamma']$ -module; (a) follows.

39.4. Let $W, I, \mathcal{X}$ be as in 39.1. Let Γ be a finite group with generators a, c and relations $a^M = 1, c^N = 1, cac^{-1} = a^k$ where M, N, k are integers ≥ 1 such that $k^N \equiv 1 \pmod{M}$. Let $\langle a \rangle$ (resp. $\langle c \rangle$) be the subgroup of Γ generated by a (resp. c). Let $\gamma \mapsto \rho_\gamma$ be a homomorphism $\Gamma \rightarrow \text{Aut}(W, I, \mathcal{X})$. We form the semidirect product $W\Gamma, W\langle a \rangle$ with W normal. Note that $W\langle a \rangle$ is a subgroup of $W\Gamma$.

Lemma 39.5. *In the setup of 39.4 let E be a simple $\bar{\mathfrak{U}}[W\Gamma]$ -module. Assume that either $cxc^{-1} = x$ for any $x \in \langle a \rangle$, or $cxc^{-1} = x^{-1}$ for any $x \in \langle a \rangle$. Let $x \in \langle a \rangle$. There exists ζ , a root of 1 in $\bar{\mathfrak{U}}$, such that $\text{tr}(wxc, E) \in \zeta \mathbf{Z}$ for any $w \in W$.*

Let $W' = W\langle a \rangle$. We can write canonically $E = \bigoplus_{t \in T} E^t$ where E^t are isotypical $\bar{\mathfrak{U}}[W']$ -modules. Now Γ acts on T by $\gamma E^t = E_{\gamma(t)}$. This action is transitive since E is simple as a $\bar{\mathfrak{U}}[W\Gamma]$ -module. If $c(t) \neq t$ for any $t \in T$ then for any $w \in W$, $wxc : E \rightarrow E$ permutes the summands E^t and no summand is stable hence $\text{tr}(wxc, E) = 0$. In this case the lemma is clear. Thus we may assume that $c(t) = t$ for some $t \in T$. Then $cE^t = E^t$ hence E^t is a $\bar{\mathfrak{U}}[W\Gamma]$ -submodule of E . Since E is simple, we have $E = E^t$. Thus, E is isotypical as a $\bar{\mathfrak{U}}[W']$ -module. Using 39.3(a) for $W\Gamma, W', \langle c \rangle$ instead of Γ, Γ', C we see that E is simple as a $\bar{\mathfrak{U}}[W']$ -module.

We can write canonically $E = \bigoplus_{h \in H} E_h$ where E_h are isotypical $\bar{\mathfrak{U}}[W]$ -modules. Now Γ acts on H by $\gamma E_h = E_{\gamma(h)}$. This action is transitive since E is simple as a $\bar{\mathfrak{U}}[W\Gamma]$ -module. The restriction of this action to $\langle a \rangle$ is also transitive since E is simple as a $\bar{\mathfrak{U}}[W']$ -module. For $h \in H$ let $\langle a \rangle_h$ be the stabilizer of h in $\langle a \rangle$. Let $W'_h = W\langle a \rangle_h \subset W'$. Then E_h is a W'_h -submodule of E and the W' -module E is induced by the W'_h -module E_h . Since E is simple as a W' -module it follows that E_h is simple as a W'_h -module. Using 39.3(a) for W'_h, W, E_h instead of Γ, Γ', E , we see that E_h is simple as a W -module.

For $h \in H$ let $\Gamma_h = \{\gamma \in \Gamma; \gamma(h) = h\}$. Then E_h is a $W\Gamma_h$ -submodule of E . Note that Γ_h is an extension of a cyclic group (the image of Γ_h under $\Gamma \rightarrow \Gamma/\langle a \rangle$) by a cyclic group (the intersection $\Gamma_h \cap \langle a \rangle$). Let $H_0 = \{h \in H; xc(h) = h\}$. For any $w \in W$ we have

$$\text{tr}(wxc, E) = \sum_{h \in H_0} \text{tr}(wxc, E_h).$$

In particular, if $H_0 = \emptyset$ then $\text{tr}(wxc, E) = 0$ so that the lemma is clear in this case. Thus we may assume that $H_0 \neq \emptyset$. Let $h \in H_0$. We can find a simple $\mathbf{Q}[W]$ -submodule E_h^0 of E_h such that $E_h = \bar{\mathfrak{U}} \otimes_{\mathbf{Q}} E_h^0$ as $\bar{\mathfrak{U}}[W]$ -modules. By Lemma 39.2 applied to W, Γ_h, E_h^0 instead of W, Γ, E we see that the $\mathbf{Q}[W]$ -structure on E_h^0 extends to a $\mathbf{Q}[W\Gamma_h]$ -module structure. (The hypotheses of that lemma are

satisfied since the $\bar{\mathfrak{U}}[W]$ -module structure on E_h extends to a $\bar{\mathfrak{U}}$ -module structure.) For $\gamma \in \Gamma_h$ let $t_\gamma \in \text{Aut}_{\mathbf{Q}}(E_h^0)$ be the action of γ in this $\mathbf{Q}[W\Gamma_h]$ -module. By extension of scalars, t_γ defines an element $\tilde{t}_\gamma \in \text{Aut}_{\mathbf{Q}}(E_h)$. For $e \in E_h, w \in W$ we have $\tilde{t}_\gamma(we) = (\gamma w \gamma^{-1})(\tilde{t}_\gamma(e))$. Hence $\gamma^{-1}(\tilde{t}_\gamma(we)) = w \gamma^{-1} \tilde{t}_\gamma(e)$. Thus $\gamma^{-1} \tilde{t}_\gamma : E_h \rightarrow E_h$ commutes with the action of W . Since E_h is simple we see that $\gamma^{-1} \tilde{t}_\gamma : E_h \rightarrow E_h$ is a scalar $\lambda_h(\gamma) \in \bar{\mathfrak{U}}^*$. Thus $\tilde{t}_\gamma = \lambda_h(\gamma) \gamma : E_h \rightarrow E_h$. Clearly $\gamma \mapsto \lambda_h(\gamma)$ is a homomorphism $\Gamma_h \rightarrow \bar{\mathfrak{U}}^*$. For $w \in W$ we have $wxc = \lambda_h(xc)^{-1} w \tilde{t}_{xc} : E_h \rightarrow E_h$ hence

$$\text{tr}(wxc, E_h) = \lambda_h(xc)^{-1} \text{tr}(w \tilde{t}_{xc}, E_h) = \lambda_{xc}^{-1} \text{tr}(wt_{xc}, E_h^0).$$

Note that $\text{tr}(wt_{xc}, E_h^0) \in \mathbf{Z}$ since E_h^0 is a $\mathbf{Q}[W\Gamma_h]$ -module. We deduce

$$\text{tr}(wxc, E) = \sum_{h \in H_0} \lambda_h(xc)^{-1} \text{tr}(wt_{xc}, E_h^0).$$

Let $H_1 = \{h \in H_0; \text{tr}(w' t_{xc}, E_h^0) \neq 0 \text{ for some } w' \in W\}$. Then, clearly,

$$\text{tr}(wxc, E) = \sum_{h \in H_1} \lambda_h(xc)^{-1} \text{tr}(wt_{xc}, E_h^0).$$

Since $\lambda_h(xc)$ is a root of 1, it is enough to verify the following statement:

If $h, h' \in H_1$ then $\lambda_h(xc) = \pm \lambda_{h'}(xc)$.

Since $\langle a \rangle$ acts transitively on H , we can find $y \in \langle a \rangle$ such that $h = yh'$. We have $yE_{h'} = E_h, y\Gamma_{h'}y^{-1} = \Gamma_h$. Also, for any $w' \in W$ we have

$$\text{tr}(w'xc, E_{h'}) = \text{tr}(yw'xcy^{-1}, E_h) = \text{tr}(yw'y^{-1}yxcy^{-1}, E_h)$$

hence

$$\lambda_{h'}(xc) \text{tr}(w'xc, E_{h'}^0) = \lambda_h(yxcy^{-1}) \text{tr}(yw'y^{-1}w'xcy^{-1}, E_h^0).$$

We take here $w' \in W$ such that $\text{tr}(w'xc, E_{h'}^0) \neq 0$. Dividing by $\text{tr}(w'xc, E_{h'}^0)$ we deduce $\lambda_{h'}(xc) \in \mathbf{Q}\lambda_h(yxcy^{-1})$. Since $\lambda_{h'}(xc), \lambda_h(yxcy^{-1})$ are roots of 1, it follows that $\lambda_{h'}(xc) = \pm \lambda_h(yxcy^{-1})$. Thus it is enough to show that $\lambda_h(xc) = \pm \lambda_h(yxcy^{-1})$ or that $\lambda_h(c^{-1}x^{-1}yxcy^{-1}) = \pm 1$ or that $\lambda_h(c^{-1}ycy^{-1}) = \pm 1$. More generally, we will verify the following statement:

If $h \in H_0, y \in \langle a \rangle, c^{-1}ycy^{-1} \in \langle a \rangle_h$ then $\lambda_h(c^{-1}ycy^{-1}) = \pm 1$.

If $cy = yc = 1$ this is obvious. Therefore we may assume that $cy' = y'^{-1}c$ for any $y' \in \langle a \rangle$.

Let $u \in \langle a \rangle_h$. Since u, xc belong to Γ_h we have $\lambda_h((xc)^{-1}uxcu^{-1}) = 1$ that is, $\lambda_h(c^{-1}ucu^{-1}) = 1$ (we use $ux = xu$). But $c^{-1}uc = u^{-1}$. Hence $\lambda_h(u^{-2}) = 1$ that is, $\lambda_h(u)^{-2} = 1$ and $\lambda_h(u) = \pm 1$. We apply this with $u = c^{-1}y_1cy_1^{-1} \in \langle a \rangle_h$. We see that $\lambda_h(c^{-1}y_1cy_1^{-1}) = \pm 1$. The lemma is proved.

39.6. Let $W, I, \mathcal{X}$ are as in 39.1. We assume that (W, I) is irreducible. Let $\tilde{I} = I \sqcup \{\omega\}$ where ω is a symbol. We define a map $\pi : \tilde{I} \rightarrow W$ as follows: $\pi(s) = s$ if $s \in I$ and $\pi(\omega)$ is the unique reflection of maximal length in $\mathcal{X}$. The restriction of π to $\tilde{I}$ is injective if $|I| \geq 2$; if $|I| = 1$, it maps both elements of $\tilde{I}$ to the unique element of I . Let Ω be the group of all permutations $\sigma : \tilde{I} \xrightarrow{\sim} \tilde{I}$ such that there exist $w \in W$ with $w\pi(\sigma)w^{-1} = \pi(\sigma(x))$ for all $x \in \tilde{I}$. Then

$$\sigma \mapsto [w_1 \mapsto ww_1w^{-1}]$$

is a homomorphism of Ω into $\text{Inn}(W)$, the group of inner automorphisms of W . Let K be a subset of $\tilde{I}$ such that $K \neq \tilde{I}$. Then π restricts to a bijection $K \xrightarrow{\sim} \pi(K)$. Let $\Omega^K = \{\sigma \in \Omega; \sigma(K) = K\}$. Let $W^{(K)}$ be the subgroup of W generated

by $\pi(K)$. From the theory of affine Weyl groups we see that $W^{(K)}$ is a (finite) Coxeter group on the generators $\pi(K)$. We have canonically $W^{(K)} = \prod_{z \in Z} W_z^{(K)}$ where $W_z^{(K)}$ is an irreducible Weyl group with set of simple reflections $K_z = K \cap W_z^{(K)}$. For $z \in Z$ we set $\mathcal{X}_z^K = R(W_z^{(K)})$ if $R(W_z^{(K)})$ is a single $W_z^{(K)}$ -conjugacy class and $\mathcal{X}_z^K = \mathcal{X} \cap W_z^{(K)}$ if $R(W_z^{(K)})$ is a union of two $W_z^{(K)}$ -conjugacy classes; in any case, $\mathcal{X}_z^K$ is a single $W_z^{(K)}$ -conjugacy class in $R(W_z^{(K)})$. (We use the following fact: if $s \in I$ is such that $s\pi(\omega)$ has order ≥ 4 then $s, \pi(\omega)$ are not conjugate under W .) Then $\mathcal{X}^K = \sqcup_{z \in Z} \mathcal{X}_z^K$ is a special subset of $R(W^{(K)})$ (see 39.1). Now the image of Ω^K under $\Omega \rightarrow \text{Inn}(W)$ (as above) is contained in the group $\text{Aut}(W^{(K)}, K, \mathcal{X}^K)$ of automorphisms of $W^{(K)}$ which preserve K and $\mathcal{X}^K$. Thus we have a homomorphism $\Omega^K \rightarrow \text{Aut}(W^{(K)}, K, \mathcal{X}^K)$. Restricting this to a subgroup C of Ω^K we obtain a homomorphism $C \rightarrow \text{Aut}(W^{(K)}, K, \mathcal{X}^K)$. Let $c \in \text{Aut}(W, I, \mathcal{X})$ be such that $c(K) = K$. We have $c^N = 1$ for some $N \geq 1$. We extend the bijection $c : I \xrightarrow{\sim} I$ (restriction of $c : W \rightarrow W$) to a bijection $c : \tilde{I} \xrightarrow{\sim} \tilde{I}$ by $c(\omega) = \omega$. We have $c(W^{(K)}) = W^{(K)}$, $c(\mathcal{X}^K) = \mathcal{X}^K$ hence c restricts to an element of $\text{Aut}(W^{(K)}, K, \mathcal{X}^K)$. For any $\sigma \in \Omega$ we define an element $c(\sigma) \in \Omega$ by $c(\sigma)(x) = \sigma(c^{-1}x)$ for $x \in \tilde{I}$. Then $c : \Omega \xrightarrow{\sim} \Omega$ preserves Ω^K . Assume that $c(C) = C$. For $\sigma \in C, w \in W^{(K)}$ we have $c(\sigma(w)) = c(\sigma)(c(w))$. Let $\langle c \rangle$ be a cyclic group of order N with generator c . On the set $W^{(K)} \times C \times \langle c \rangle$ we have a group structure

$$(w, \sigma, c^n)(w', \sigma', c^{n'}) = (w\sigma(c^n(w')), \sigma c^n(\sigma'), c^{n+n'}).$$

This is the semidirect product $W^{(K)}\Gamma$ of $W^{(K)}$ with the group $\Gamma = C \times \langle c \rangle$ with group structure $(\sigma, c^n)(\sigma', c^{n'}) = (\sigma c^n(\sigma'), c^{n+n'})$.

Proposition 39.7. *In the setup of 39.6 let E be a simple $\bar{\mathfrak{U}}[W^{(K)}\Gamma]$ -module. Let $\gamma \in \Gamma$. There exists ζ , a root of 1 in $\bar{\mathfrak{U}}$, such that $\text{tr}(w\gamma, E) \in \zeta \mathbf{Z}$ for any $w \in W^{(K)}$.*

From the theory of affine Weyl groups it is known that one of the following holds:

- (i) Ω is cyclic and $c(\sigma) = \sigma$ for all $\sigma \in \Omega$.
- (ii) Ω is cyclic and $c(\sigma) = \sigma^{-1}$ for all $\sigma \in \Omega$,
- (iii) $\Omega \cong \mathbf{Z}/2 \times \mathbf{Z}/2$.

Moreover, $C \subset \Omega$ (compatibly with the action of c). In cases (i),(ii), C is cyclic and the assumptions of 39.5 are satisfied (with $W^{(K)}$ instead of W). Hence the result follows from 39.5.

In the remainder of the proof we assume that we are in case (iii). If C is cyclic then $c(\sigma) = \sigma = \sigma^{-1}$ for all $\sigma \in C$. The result follows again from 39.5. Hence we may assume that C is not cyclic so that $C = \Omega$. Let n be the order of $c : C \rightarrow C$. Then $n \in \{1, 2, 3\}$ and n divides N . Note that $c^{N/n}$ is in the centre of Γ . Tensoring E by a suitable one dimensional representation (which is trivial on $W^{(K)}C$) we may assume that $c^{N/n}$ acts trivially on E hence E factors through the quotient of $W^{(K)}\Gamma$ by the subgroup generated by $c^{N/n}$. Hence we may assume that $N = n$. If $n = 1$ then $\Gamma \cong \mathbf{Z}/2 \times \mathbf{Z}/2$ is a group as in 39.5 and the result follows from 39.5.

If $n = 2$ then Γ is a dihedral group of order 8 which is again a group as in 39.5 (an extension of $\mathbf{Z}/2$ by $\mathbf{Z}/4$) and the result follows from 39.5. If $n = 3$ then W must be of type D_4 , $W^{(K)}$ is an elementary abelian 2-group, E is one dimensional and there exists a homomorphism $\mu : W^{(K)}\Gamma \rightarrow \bar{\mathfrak{U}}^*$ such that $\text{tr}(w\gamma', E) = \mu(w\gamma')$ for all $w \in W^{(K)}$, $\gamma' \in \Gamma$. Then $\zeta = \mu(\gamma)$ is a root of 1. For any $w \in W^{(K)}$ we have $\mu(w\gamma) = \zeta\mu(w)$ and $\mu(w) = \pm 1$, since $w^2 = 1$. Thus, $\text{tr}(w\gamma, E) = \pm\zeta$. The proposition is proved.

Corollary 39.8. *Assume that G, D, A, u are as in 35.22 and that G^0/Z_{G^0} is simple. Then $b_{A,u}^v \in \eta\mathbf{Q}$ for some η , a root of 1 (with $b_{A,u}^v$ as in 34.19.)*

This follows from 35.22 and 39.7.

REFERENCES

- [Al] D. Alvis, *The duality operation in the character ring of a finite Chevalley group*, Bull. Amer. Math. Soc. **1** (1979), 907-911.
- [BBD] A. Beilinson, J. Bernstein, P. Deligne, *Faisceaux pervers*, Astérisque **100** (1982).
- [Cu] C.W. Curtis, *Truncation and duality in the character ring of a finite group of Lie type*, J. Algebra **62** (1980), 320-332.
- [De] P. Deligne, *La conjecture de Weil, II*, Publ.Math. I.H.E.S. **52** (1980), 137-252.
- [Ka] N. Kawanaka, *Fourier transforms of nilpotently supported invariant functions on a simple Lie algebra over a finite field*, Invent.Math. **69** (1982), 411-435.
- [L14] G. Lusztig, *Characters of reductive groups over a finite field*, Ann. Math. Studies 107, Princeton U.Press, 1984.
- [L3] G. Lusztig, *Character sheaves, I*, Adv. Math. **56** (1985), 193-237; II **57** (1985), 226-265; III **57** (1985), 266-315; IV **59** (1986), 1-63; V **61** (1986), 103-155.
- [L9] G. Lusztig, *Character sheaves on disconnected groups, I*, Represent. Th. (electronic) **7** (2003), 374-403; II **8** (2004), 72-124; III **8** (2004), 125-144; IV **8** (2004), 145-178; Errata **8** (2004), 179-179; V **8** (2004), 346-376; VI **8** (2004), 377-413; VII **9** (2005), 209-266.
- [L10] G. Lusztig, *Parabolic character sheaves, I*, Moscow Math.J. **4** (2004), 153-179.

DEPARTMENT OF MATHEMATICS, M.I.T., CAMBRIDGE, MA 02139